



Dr. Sandra L. Anagnostakis
Department of Plant Pathology and Ecology
The Connecticut Agricultural Experiment Station
123 Huntington Street, P. O. Box 1106
New Haven, CT 06504

Phone: (203) 974-8498

Fax: (203) 974-8502

E-mail: Sandra.Anagnostakis@ct.gov

Website: www.ct.gov/caes

Founded in 1875

Putting science to work for society

GROWING CHESTNUT TREES

Why grow chestnuts? A few acres of land can yield nuts for your own enjoyment or for sale at the side of the road or to your local market. Chestnuts are a good food, as shown in this table:

<u>Species</u>	<u>Fiber %</u>	<u>Protein %</u>	<u>Fat %</u>	<u>Carbohydrates %</u>	<u>Calories per ounce</u>
Chinese	14	8	2	65	66
European	14	6	4	66	57
Japanese	14	8	0.4	90	114
American	19	10	10	40	70

This is a guide for choosing the right chestnut species, choosing the right planting site, and anticipating the care that is needed. For a few trees for your own use, you would do well to get two or three cultivars and plant them near each other (within 100 feet) for optimal cross-pollination. If you are starting an orchard for commercial production, you would be better off with a lot of trees of a few cultivars and a couple of seedling trees to provide pollen. The advantage of a uniform orchard is that the nuts all ripen at the same time, and will be fairly uniform in size and appearance (something consumers seem to think is important). The other point to remember for selling the nuts is that “bigger is worth more” even though smaller nuts frequently taste better!

All species of chestnuts can pollinate each other, but hybrids often produce no pollen (are “male sterile”) and this must be determined when planning your orchard. Chestnut Blight Disease must be considered in the eastern U.S., Ink Disease (caused by *Phytophthora*) may be a problem in the south or in areas with poor drainage, Chestnut Gall Wasp is slowly moving north from Georgia, Alabama, North and South Carolina, and Tennessee and was recently found in Ohio, and drought tolerance will make a big difference in the west.

The initial investment will include purchase of trees, tree shelters to protect them, and installation of an irrigation system.

“Named cultivars” are clones of trees whose characteristics are known. Since they are identical, and chestnuts will not self-pollinate, two or more cultivars (or some seedlings, which are all different) must be planted to provide pollen for each other. Cultivars are grafted onto compatible rootstocks, and they usually cost about \$15 each (see “sources” list appended). The choice of

cultivar must be made for your own environmental conditions and taste, and some guidelines can be found in our list of cultivars (e-mail for a copy). Information on nut size has not been systematically collected in all of the U.S. growing regions for cultivars, so the following is only to provide a general range of the sizes available:

<u>Cultivar</u>	<u>Number of nuts per pound*</u>
'Okei'	12-15
'Colossal'	14-18
'Bouche de Betizac'	15
'Willamette'	18-22
'Qing'	20
'Sleeping Giant'	28-34

*From J. M. Nave. 1998. Large Fruited Chestnuts Grown in North America, 89th Annual Report of the Northern Nut Growers Association, pages 42-82.

Tree shelters can be made of wire or plastic, or can be as simple as piled brush to protect the trees from deer. The plastic tubes cost about \$3.00 each and are easy to install, but must be lifted each fall to allow the trees to harden off properly. They frequently harbor wasps, and provide a nice environment for rodents who then eat the tree bark. A better shelter may be a sturdy enclosure of hardware cloth (expanded metal) about 3 feet in diameter.

An irrigation system is essential for a commercial orchard, even if you live in a region where rainfall is fairly dependable.

Plant your trees in well-drained sandy loam, better drained than apples require. Sites with clay may be tolerated if there is good surface drainage (slope). Chestnuts need a soil pH of 5.5 to 6.5 and absolutely won't tolerate limestone soils. If your soil pH is much higher than this, consider another crop.

Space the trees at 40 feet apart, or start at 25 feet apart and remove trees, as they mature, to 35 feet apart.

Fertilize (SPRING ONLY) with one pound of 5-10-5 or 10-10-10 per year of tree age, or one pound per inch of tree trunk diameter.

Pruning can be done when it is hot and dry, in early summer (never later than the middle of July). This reduces the chance of infection. Pruning in late winter while trees are dormant is also possible.

The area around the trees should be kept mowed, and a circle about three feet wider than the diameter of the trees should be kept weed-free. Mulch can be applied to help with water retention, but should not be deeper than two inches, and should not be up against the trunks (allowing easy access by rodents in the winter).

The value of the nuts is directly related to the size, but is usually at least \$1.50 per pound wholesale and up to \$5.00 per pound retail. Yield will start to be significant after the trees are 10 years old, and yields have been reported of from 14 pounds per tree to 138 pounds per tree.

The traditional method of harvest is to allow the nuts to fall to the ground, and employ people to pick them up. For a few nuts for your own table this is not a bad system, but commercial

orchards need more efficient methods. Nets can be suspended under the bottom branches of the trees to collect the nuts (they should be removed from the nets every day), or some of the marketed ground-collection systems can be used (see advertisements in the Northern Nut Growers Newsletter).

The most serious problem for all U.S. chestnut growers is the presence of chestnut weevils that lay their eggs in the ripening nuts. When the nuts fall to the ground, the eggs hatch, and the larvae eat their way out of the nuts and burrow into the ground. They stay there until the following year (Large Chestnut Weevil) or the second year (Small Chestnut Weevil) when they emerge to mate and lay more eggs. Only the adult weevils are controllable with insecticides (Sevin is registered for this use), and care must be taken to spray the burs weekly for the last month before nuts are released. An “organic” control that has proved reasonably effective is the deployment of chickens or Guinea fowl under the trees to eat the larvae and pupae of the weevils. After several years of this control, few weevils will remain. As soon as nuts are collected they can be treated in hot water (120 °F = 49 °C) for 20 minutes to kill weevil eggs and larvae but not the nuts. Then the nuts should be refrigerated. Keep them moist, in plastic bags, in the refrigerator (over the winter) if you are planning to plant them, or a little dryer for eating. Nuts can also be par-boiled as soon as they are harvested and stored (frozen) in plastic bags until needed. This kills the eggs (or very small larvae) before they can damage the nuts but also kills the nuts.

Marketing chestnuts is easy if you already have a farm stand for selling produce, but if you have to sell them to local food stores, you will have to work a little harder. Nuts should never be allowed to dry out, and markets must be convinced to display and store them in plastic bags in refrigerated cases.

The future of the chestnut industry in this country will depend on the development of better weevil control methods, better harvesting techniques, and development of “ready for use products.” Peeled, dried nuts, or chips that can be added directly to foods being prepared will allow cooks to experiment, and may interest large, prepared-food manufacturers to include chestnuts in their products.

SOURCES OF CHESTNUT TREES

American Chestnut Cooperators Foundation

<http://ipm.ppws.vt.edu/griffin/accf.html>

seedlings from American chestnuts with some resistance to chestnut blight

Burnt Ridge Nursery 360-985-2873
mail@burntridgenursery.com
www.burntridgenursery.com
432 Burnt Ridge Rd., Onalaska, WA 98570
'Colossal', 'Nevada', 'Layeroka', 'Skioka',
'Skookum', 'Maraval', grafted cultivars: Chinese chestnuts, American chestnuts and
chinquapins, European chestnuts: seedlings available as well

Chestnut Hill Nursery 800-669-2067
www.ChestnutHillTreeFarm.com
chnuts@msn.com
15105 NW 94th Ave., Alachua, Florida 32615
seedlings

Empire Chestnut Company 330-627-3181
www.EmpireChestnut.com
empire@eohio.net
3276 Empire Rd SW, Carrollton, OH 44615-9515
'Amy', 'Gideon', 'Eaton', 'Peach', 'Sleeping Giant', 'Paragon', 'Qing', and
archived 'AU-Homestead', 'Carr', 'Chandler', 'Crane', 'Kohr', 'Lockwood', 'Norris',
'Nanking', 'Orrin', 'Smith' and others, Chinese seedlings, seedling *Castanea henryi*,
Chinquapin seedlings, grafted cultivars

England's Orchard and Nursery 606-965-2228
www.nutttrees.net
nutttrees@prtcnet.org
316 S.R. 2004, Mckee, KY 40447-9616
grafted and seedling trees

Forrest Keeling Nursery 800-356-2401
info@fknursery.com
www.fknursery.com
88 Forrest Keeling Lane, Elsberry, Missouri 63343
grafted trees of 'Dunston', 'Eaton', 'Gideon', 'Homestead', 'Improved Chinese
Chestnut', 'Peach', 'Sleeping Giant', 'Colossal'

Grimo Nut Nursery 905-934-6887
979 Lakeshore Rd., R.R. 3, Niagara-on-the-Lake, Ontario LOS 1JO, Canada
nutttrees@grimonut.com
www.grimonut.com
numbered types and seedlings

Lawyer Nursery 406-826-3881, fax. 406-826-5700

www.lawyernursery.com

950 Highway 200 West, Plains, Montana 59859

seed of Chinese and American chestnut

seedlings, 1 year old

Nolin River Nut Tree Nursery 270-369-8551

www.nolinnursery.com

797 Port Wooden Rd., Upton, Kentucky 42784

'Armstrong' 'Au Leader' 'Crane' 'Eaton River' 'Ford's Sweet' 'Ford's Tall'

'Meiling' 'Mendes' 'Mossbarger' 'Orrin' 'Qing' 'Sleeping Giant'

Northwoods Wholesale Nursery 503-651-3737

28696 S. Cramer, Molalla, Oregon 97038

seedling Chinese and Japanese X European chestnuts

WHOLESALE ONLY

Nursery Street Greenhouses 616-624-1806 or 616-624-6054

ptibyrne@aol.com

61591 30th Street, Lawton, Michigan 49065

rooted cuttings of 'Colossal' 'Nevada' 'Skookum' several Grimo clones, several

Higgins clones, and several Adkins clones

Oikos Tree Crops 616-624-6233, fax. 616-624-4019

oak24@aol.com

www.oikostreecrops.com/

P.O. Box 19425, Kalamazoo, Michigan 49019-0425

seedlings of Chinese hybrids (Douglas) and others

Plantra Tree Shelters (800) 951-3806

www.plantra.com

2508 Northland Drive

Mendota Heights, MN 55120

Tree shelters of several kinds

Red Fern Farm 319-729-5905

RedFernFarm@lisco.com

www.RedFernFarm.com

13882 "P" Ave., Wapello, IA 52653-9449

Chinese chestnut seedlings

Rhora's Nut Farm and Nursery 905-899-3508
32983 Wills Rd., RR #1
Wainfleet, Ontario, Canada
seedling chestnuts

St. Lawrence Nurseries 315-265-6739
www.sln.potsdam.ny.us
trees@sln.potsdam.ny.us
325 State Hwy 345, Potsdam, NY 13676
seedling chestnut trees sometimes available

Stark Brothers Nursery 800-325-4180
www.starkbros.com
Louisiana, MO 63353
'Colossal' and 'Starks Choice', seedling trees

Sticks and Stones Nursery 931-668-2409
gclendenon@blomand.net
www.SticksAndStonesNursery.com
526 Center Hill Rd., McMinnville, TN 37110
"Sweet Hart" chestnut seedlings, "Mastmaker" chestnut seedlings, Allegheny chinquapin seedlings

Superior Trees, Inc., Lee Nursery 904-971-5159
P.O. Box 9325, U.S. 90 East, Lee, Florida 32059
Allegheny chinquapin seedlings, price varies by number ordered

Wexford Soil and Water Conservation District 231-775-7422
7192 East 34 Rd., Cadillac, Michigan 49601
American chestnut seedlings

Woodlanders, Inc. 803-648-7522
www.woodlanders.net
1128 Colleton Ave., Aiken, SC 29801
seedling Chinese chestnuts, seedling chinquapins, of several species
(including type "Herschel").

January 2013 (revised)