

Adolescent ASAM Criteria 2001 <i>Cliff Notes</i>	Outpatient Treatment	Intensive Outpatient Treatment (IOP/SOPS)	Partial Hospitalization
	<i>Level I</i> <i>*Generally &lt; 6 hours/wk</i>	<i>Level II.1</i> <i>*Generally 6+ hours/wk</i>	<i>Level II.5</i> <i>*Generally 20+ hours/wk</i>
<b>Dimension 1</b> Alcohol Intoxication and/or Withdrawal Potential	<b>None</b>	<b>Mild OR</b> is at risk of withdrawal	<b>Mild OR</b> is at risk of withdrawal
<b>Dimension 2</b> Biomedical Conditions and Complications	<b>None or Mild</b>	<b>None or Mild</b> Distracts from treatment in a lower level of care.	<b>None or Mild</b> Distracts from treatment in a lower level of care.
<b>Dimension 3</b> Emotional, Behavioral or Cognitive Conditions and Complications	Status is characterized by <b>ALL</b> <b>of the following:</b> <b>a, b, c, d, &amp; e</b>	Status features <b>One or More</b> <b>of the following:</b> <b>a, b, c, d, &amp; e</b>	Status features <b>One or More</b> <b>of the following:</b> <b>a, b, c, d, &amp; e</b>
a) Dangerousness/Lethality	<b>None</b>	<b>Mild</b> Is safe between sessions	<b>Mild</b> Is safe overnight
b) Interference With Addiction Recovery Efforts	<b>Mild</b>	<b>Mild</b> This level of care required to support treatment engagement	<b>Moderate</b> This level of care required to support treatment engagement
c) Social Functioning	<b>None to Mild</b> Impairment	<b>Mild to Moderate</b> but can sustain responsibilities	<b>Moderate</b> but can sustain responsibilities
d) Ability For Self-Care <i>*Activities of Daily Living</i>	<b>Mild</b> but there is significant risk of deterioration	<b>Mild to Moderate</b> and requires frequent monitoring or interventions	<b>Moderate</b> and requires almost daily monitoring or interventions
e) Course Of Illness <i>*What can we tell from their history, patient has a pattern of...</i>	<b>Mild</b> (imminent) risk which predicts a need for some monitoring or interventions	<b>Moderate</b> History combined with current situation predicts need for frequent monitoring or interventions	<b>Moderate</b> History combined with current situation predicts need for almost daily monitoring or interventions
<b>Dimension 4</b> Readiness to Change	Willing to engage in treatment and is at least contemplating change, but needs motivating and monitoring strategies.	<b>Moderate</b> Requires close monitoring and support several times/wk to promote progress because of variable treatment engagement or a lack of recognition of the need for assistance.	<b>Moderate</b> Requires almost daily structure to promote progress because of poor treatment engagement, escalating use and impairment or lack of recognition of the role of substances in their present problems.
<b>Dimension 5</b> Relapse, Continued Use or Continued Problem Potential	Needs limited support to maintain abstinence or control use and to pursue recovery goals.	<b>Moderate</b> Needs close monitoring & support because of a significant risk of relapse or continued use & deterioration in level of functioning. Poor relapse prevention skills.	<b>Moderate</b> Needs almost daily monitoring & support because of a high risk of relapse or continued use & deterioration in level of functioning. Minimal relapse prevention skills.

Adolescent ASAM Criteria 2001 <i>Cliff Notes</i>	Outpatient Treatment	Intensive Outpatient Treatment (IOP/SOPS)	Partial Hospitalization
	<i>Level I</i> <i>*Generally &lt; 6 hours/wk</i>	<i>Level II.1</i> <i>*Generally 6+ hours/wk</i>	<i>Level II.5</i> <i>*Generally 20+ hours/wk</i>
<b>Dimension 6</b> Recovery Environment	Family/Environment can support recovery with limited assistance.	<b>Moderate</b> Environment is impeding their recovery. Requires close monitoring and support to overcome that barrier.	<b>Moderate</b> Environment renders recovery unlikely without almost daily monitoring and support or frequent relief from his or her home environment.

\*Summary based on ASAM Patient Placement Criteria, Second Edition-Revised, (Crosswalk) Page 191-193

August 2001