

Adolescent ASAM Criteria 2001 Cliff Notes	Clinical Managed Medium Intensity Residential	Medical Monitored High Intensity Residential Inpatient	Medically Managed Intensive Inpatient
	<i>Therapeutic Community Residential Rehab</i>	<i>IP Rehab IP Residential</i>	<i>Inpatient Detox 23 Hour Bed</i>
	<i>Level III.5</i>	<i>Level III.7</i> <i>*Require (1) from D1-3 &amp; D4-6 Be Met</i>	<i>Level IV</i>
<b>Dimension 1</b> Alcohol Intoxication and/or Withdrawal Potential	<b>Mild or Moderate</b> withdrawal (or at risk of) <i>but does not need</i> pharmacological management or frequent medical/nursing monitoring	<b>Moderate to Severe</b> Withdrawal or Risk but is manageable at III.7	<b>Severe</b> Withdrawal or Risk Requires intensive active medical management
<b>Dimension 2</b> Biomedical Conditions and Complications	<b>None or Mild</b> Receiving medical monitoring as needed	<b>Moderate</b> Requires medical monitoring but not intensive treatment	<b>Severe</b> Requires 24hr medical/nursing care
<b>Dimension 3</b> Emotional, Behavioral or Cognitive Conditions and Complications	Status features <b>One or More of the following:</b> <b>a, b, c, d &amp; e</b>	Status features <b>One or More of the following:</b> <b>a, b, c, d &amp; e</b>	Status features <b>One or More of the following:</b> <b>a, b, c, d &amp; e</b>
a) Dangerous/Lethality	<b>Moderate</b> but stable risk of harm, and thus needs medium-intensity, 24hr monitoring or treatment for safety	<b>Moderate</b> Needs high-intensity 24hr monitoring or treatment, or secure containment, for safety	<b>Severe</b>
b) Interference with Addiction Recovery Efforts	<b>Moderate to Severe</b> Requires medium-intensity residential treatment to support engagement	<b>Severe</b> Requires high-intensity residential treatment to support engagement	<b>Very Severe</b> Almost overwhelming and renders them incapable of participating in treatment at a less intensive level of care
c) Social Functioning	<b>Moderate to Severe</b> Cannot be managed at a less intensive level of care	<b>Severe</b> Cannot be managed at a less intensive level of care	<b>Very Severe</b> Dangerous impairment and requires frequent medical/nursing interventions
d) Ability for Self-Care	<b>Moderate to Severe</b> Requires 24hr supervision and medium-intensity staff assistance	<b>Severe</b> Requires 24hr supervision and high-intensity staff assistance	<b>Very Severe</b> Requires frequent medical/nursing interventions
<b>Adolescent ASAM Criteria 2001*Cliff Notes</b>	<i>Level III.5 Residential Rehab</i>	<i>Level III.7 IP Rehab/IP Residential</i>	<i>Level IV IP Detox/23 Hour Bed</i>

<p>e) Course of Illness  <i>*What can we tell from their history, patient has a pattern of...</i></p>	<p><b>Moderate</b>  History combined with current situation predicts destabilization without medium-intensity residential treatment</p>	<p><b>Moderate to Severe</b>  History combined with current situation predicts destabilization without high-intensity residential treatment</p>	<p><b>Severe</b>  History combined with current situation predicts destabilization without inpatient medical management</p>
<p><b>Dimension 4</b>  Readiness to Change</p>	<p><b>Moderate</b>  Needs intensive motivating strategies in a 24hr structured program to address minimal treatment engagement or opposition to treatment, and/or their lack of recognition of current severe impairment</p>	<p><b>Moderate or Severe</b>  Needs high-intensity motivating strategies in a 24hr medically monitored program to address their lack of treatment engagement associated with a biomedical, emotional or behavioral condition; or because they are actively opposed to treatment requiring confinement; or they need high-intensity case mgmt, to create linkages that would support outpatient treatment</p>	<p>Problems in Dimension 4 do not qualify for Level IV services</p>
<p><b>Dimension 5</b>  Relapse, Continued Use or Continued Problem Potential</p>	<p><b>Moderate or Severe</b>  Unable to control use and avoid serious impairment without a 24hr structured program because he/she is unable to overcome environmental triggers or cravings; or has insufficient supervision between encounters at a less intensive level of care; or has a high chronicity and/or poor response to treatment</p>	<p><b>Severe</b>  Unable to interrupt a high severity or high frequency pattern of use and avoid dangerous consequences without high-intensity 24hr interventions (because of an emotional, behavioral or cognitive condition, severe impulse control problems, withdrawal symptoms, and the like)</p>	<p>Problems in Dimension 5 do not qualify for Level IV services</p>
<p><b>Dimension 6</b>  Recovery Environment</p>	<p><b>Moderate or Severe</b>  Environment is dangerous to his/her recovery, so that he/she requires residential treatment to promote recovery goals or for protection.</p>	<p><b>Severe</b>  Environment is dangerous to their recovery, and they require residential treatment to promote recovery goals or for protection, and to help them establish a successful transition to a less intensive level of care.</p>	<p>Problems in Dimension 6 do not qualify for Level IV series</p>
<p>*Summary based on ASAM Patient Placement Criteria, Second Edition-Revised, (Crosswalk) Page 194-197 August 2001</p>			