

Number and percentage of persons receiving HIV care^e between 01/01/2015 through 12/31/2015 among persons who were >=13 years old on 12/31/2014, resided in Connecticut (based on most recent residence), diagnosed with HIV infection through 12/31/2014 and living with HIV on 12/31/2015, by selected characteristics- based on HIV surveillance data reported through 12/23/2016.

| Characteristics | No. of persons diagnosed with HIV infection through 12/31/2014 and living with HIV on 12/31/2015 (overall population) ^h | No. of persons who have >=1 care visit ^f between 01/01/2015 through 12/31/2015 | No. of persons who have >=2 care visits between 01/01/2015 through 12/31/2015, at least 3 months apart ^g | % of persons who have >=2 care visits between 01/01/2015 through 12/31/2015, at least 3 months apart among the overall population | % of persons who have >=2 care visits between 01/01/2015 through 12/31/2015, at least 3 months apart among persons have >=1 care visit between 01/01/2015 through 12/31/2015 |
|--|--|---|---|---|--|
| Sex | | | | | |
| Male | 6599 | 5132 | 4134 | 62.65 | 80.55 |
| Female | 3843 | 3104 | 2459 | 63.99 | 79.22 |
| Age on 12/31/2014 | | | | | |
| 13-24 | 301 | 235 | 181 | 60.13 | 77.02 |
| 25-34 | 1016 | 784 | 575 | 56.59 | 73.34 |
| 35-44 | 1963 | 1499 | 1176 | 59.91 | 78.45 |
| 45-54 | 3679 | 2947 | 2358 | 64.09 | 80.01 |
| 55-64 | 2732 | 2181 | 1803 | 66 | 82.67 |
| >=65 | 751 | 590 | 500 | 66.58 | 84.75 |
| Race/ethnicity | | | | | |
| American Indian/Alaska Native | 13 | 9 | 8 | 61.54 | 88.89 |
| Asian ^a | 89 | 59 | 46 | 51.69 | 77.97 |
| Black/African American | 3539 | 2828 | 2255 | 63.72 | 79.74 |
| Hispanic/Latino | 3554 | 2718 | 2236 | 62.92 | 82.27 |
| Native Hawaiian/Other Pacific Islander | 5 | 3 | 2 | 40 | 66.67 |
| White | 3124 | 2515 | 1973 | 63.16 | 78.45 |
| Multiple races | 118 | 104 | 73 | 61.86 | 70.19 |
| Transmission category | | | | | |
| Male-to-male sexual contact (MSM) | 2709 | 2205 | 1689 | 62.35 | 76.6 |
| Injection drug use (IDU) | 3177 | 2468 | 2073 | 65.25 | 84 |
| MSM and IDU | 218 | 179 | 151 | 69.27 | 84.36 |
| Heterosexual contact | 2976 | 2487 | 1974 | 66.33 | 79.37 |
| NIR/NRR (unknown) | 1164 | 753 | 578 | 49.66 | 76.76 |
| Other ^d | 198 | 144 | 128 | 64.65 | 88.89 |
| MSM | | | | | |
| American Indian/Alaska Native | 5 | 4 | 3 | 60 | 75 |
| Asian ^a | 37 | 29 | 20 | 54.05 | 68.97 |
| Black/African American | 567 | 460 | 346 | 61.02 | 75.22 |
| Hispanic/Latino | 662 | 525 | 422 | 63.75 | 80.38 |
| Native Hawaiian/Other Pacific Islander | 5 | 3 | 2 | 40 | 66.67 |
| White | 1406 | 1159 | 878 | 62.45 | 75.75 |
| Multiple races | 27 | 25 | 18 | 66.67 | 72 |
| Subtotal | 2709 | 2205 | 1689 | 62.35 | 76.6 |
| Injection drug use (male) | | | | | |
| American Indian/Alaska Native | 1 | 0 | 0 | 0 | . |
| Asian ^a | 6 | 3 | 3 | 50 | 100 |
| Black/African American | 692 | 554 | 473 | 68.35 | 85.38 |
| Hispanic/Latino | 885 | 622 | 532 | 60.11 | 85.53 |
| White | 392 | 309 | 264 | 67.35 | 85.44 |
| Multiple races | 18 | 17 | 14 | 77.78 | 82.35 |
| Subtotal | 1994 | 1505 | 1286 | 64.49 | 85.45 |
| Injection drug use (female) | | | | | |
| American Indian/Alaska Native | 1 | 1 | 1 | 100 | 100 |
| Asian ^a | 2 | 1 | 1 | 50 | 100 |
| Black/African American | 401 | 332 | 266 | 66.33 | 80.12 |
| Hispanic/Latino | 443 | 352 | 293 | 66.14 | 83.24 |
| White | 313 | 255 | 211 | 67.41 | 82.75 |
| Multiple races | 23 | 22 | 15 | 65.22 | 68.18 |
| Subtotal | 1183 | 963 | 787 | 66.53 | 81.72 |
| Heterosexual contact (male) | | | | | |
| American Indian/Alaska Native | 1 | 1 | 1 | 100 | 100 |
| Asian ^a | 9 | 6 | 6 | 66.67 | 100 |
| Black/African American | 421 | 346 | 283 | 67.22 | 81.79 |
| Hispanic/Latino | 288 | 231 | 199 | 69.1 | 86.15 |
| White | 190 | 158 | 126 | 66.32 | 79.75 |
| Multiple races | 9 | 8 | 6 | 66.67 | 75 |
| Subtotal | 918 | 750 | 621 | 67.65 | 82.8 |
| Heterosexual contact (female) | | | | | |
| American Indian/Alaska Native | 3 | 1 | 1 | 33.33 | 100 |
| Asian ^a | 10 | 9 | 6 | 60 | 66.67 |
| Black/African American | 870 | 722 | 563 | 64.71 | 77.98 |
| Hispanic/Latino | 791 | 682 | 531 | 67.13 | 77.86 |
| White | 361 | 304 | 241 | 66.76 | 79.28 |
| Multiple races | 23 | 19 | 11 | 47.83 | 57.89 |
| Subtotal | 2058 | 1737 | 1353 | 65.74 | 77.89 |
| Total | 10442 | 8236 | 6593 | 63.14 | 80.05 |

^aIncludes Asian/Pacific Islander legacy cases.

^bIncludes hemophilia, blood transfusion, and persons (171 cases) who were exposed perinatally but were aged >13 at most recent year.

^cPersons who have at least one CD4 or viral load or HIV-1 genotype test are considered as receiving HIV care.

^dTests that have been done in the same month are considered as one care visit, even two tests that one is CD4 test and the other one is viral load (or genotype) test are still considered as one care visit. ^eTwo care visits should be 3 months apart.

^fThe overall population is overestimated because cases are only followed up for 11 months after 12/31/2015. CDC suggests that every case should be followed up at least 18 months to collect death certificate information.