



STATE OF CONNECTICUT

DEPARTMENT OF PUBLIC HEALTH

BLAST FAX 14-

TO: Hospital Administrators/Nursing Home Administrators/Nursing Home Medical Directors/Infection Control Practitioners/Home Health Agency Administrators/Assisted Living Services Agencies/Residential Care Homes

FROM: Barbara Cass, R.N., Section Chief *BSL*
Facility Licensing and Investigations Section
410 Capitol Avenue
Hartford, Connecticut, 06134

DATE: May 16, 2014

SUBJECT: Use of Insulin Pens in Healthcare Facilities

On May 18, 2012, Survey and Certification (S+C) Letter 12-30 was released by the Centers for Medicare and Medicaid Services reinforcing that Insulin Pens are indicated for single patient use only. It has recently been reported to the Department that single patient insulin pens had been used for administration of insulin at a Connecticut healthcare facility inconsistent with the manufacturer's intent.

Please see the attached S+C Letter for further guidance and direction. Additionally, the CDC Clinical reminder for the use of Insulin pens is also attached.

Should you have any questions, please contact myself or Donna Ortelle, R.N., M.S.N., Public Health Services Manager at 860-509-7407.



Office of Clinical Standards and Quality/Survey & Certification Group

Ref: S&C: 12-30-ALL

DATE: May 18, 2012
TO: State Survey Agency Directors
FROM: Director
Survey and Certification Group
SUBJECT: Use of Insulin Pens in Health Care Facilities

Memorandum Summary

Insulin Pen devices: The Centers for Medicare & Medicaid Services (CMS) has recently received reports of use of insulin pens for more than one patient, with at least one 2011 episode resulting in the need for post-exposure patient notification. These reports indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients. **Insulin pens are meant for use by a single patient only.** Each patient/resident must have his/her own. Sharing of insulin pens is essentially the same as sharing needles or syringes, and must be cited, consistent with the applicable provider/supplier specific survey guidance, in the same manner as re-use of needles or syringes.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times by a single patient/resident, using a new needle for each injection. Insulin pens must never be used for more than one patient/resident. Regurgitation of blood into the insulin cartridge after injection will create a risk of bloodborne pathogen transmission if the pen is used for more than one patient/resident, even when the needle is changed [1]. A previous memo (10-28-NH), dated August 27, 2010, similarly identified that point of care testing devices must not be shared between residents because of the risk of bloodborne pathogen transmission.

In 2009, in response to reports of improper use of insulin pens in hospitals, the Food and Drug Administration (FDA) issued an alert for healthcare professionals reminding them that insulin pens are meant for use by a single patient only and are not to be shared between patients [2]. Despite this alert, patients continue to be placed at risk of bloodborne pathogen exposure through inappropriate use of insulin pens for more than one patient, including an incident in 2011 that required notification of more than 2,000 patients [3]. These events indicate that some healthcare personnel may be unaware of the risk this unsafe practice poses to patients/residents.

Discussion

Any provider or supplier using insulin pens should review the following recommendations of the FDA to prevent transmission of bloodborne infections in the patients/residents under their care.

- Insulin pens containing multiple doses of insulin are meant for single patient/resident use only, and must never be used for more than one person, even when the needle is changed.
- Insulin pens must be clearly labeled with patient/resident's name or other identifiers to verify that the correct pen is used on the correct patient/resident.
- Healthcare facilities should review their policies and procedures and educate their staff regarding safe use of insulin pens.

Reuse of insulin pens is similar to reusing needles or syringes for more than one patient/resident and must direct the surveyor to focus on the overall infection control practices in the facility. The facility plan of correction should include notification of the local health department or state epidemiologist for determination of the need for post-exposure follow-up of patients and residents.

Effective Date: Immediately. Please ensure that state and RO surveyors are incorporating this information into their survey practices.

Training: The information must be shared with all survey and certification staff, surveyors, managers, and the State and CMS Regional Office training coordinators.

Additional Resource Material

The CDC has updated their website reference material www.cdc.gov/injectionsafety and issued a clinical reminder accessible a <http://www.cdc.gov/injectionsafety/clinical-reminders/insulin-pens.html>

References

1. Sonoki K, Yoshinari M, Iwase M, Tashiro K, Iino K, Wakisaka M, Fujishima M. Regurgitation of blood into insulin cartridges in the pen-like injectors. *Diabetes Care*. 2001;24(3):603-4.
2. Information for healthcare professionals: risk of transmission of blood-borne pathogens from shared use of insulin pens (2009). *U.S. Food and Drug Administration Postmarket Drug Safety Information for Patients and Providers*. Accessed November 14, 2011 from <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/DrugSafetyInformationforHealthcareProfessionals/ucm133352.htm>
3. Important Patient Safety Notification (2011). *Dean Clinic*. Accessed November 14, 2011 from <http://www.deancare.com/about-dean/news/2011/important-patient-safety-notification/>

Page 3- State Survey Agency Directors

Questions may be sent to Karen Hoffmann at Karen.Hoffmann@cms.hhs.gov or phone (919)-622-4811.

/s/
Thomas E. Hamilton

cc: Survey and Certification Regional Office Management

CDC CLINICAL REMINDER


Insulin Pens Must Never Be Used for More than One Person

Summary

The Centers for Disease Control and Prevention (CDC) has become increasingly aware of reports of improper use of insulin pens, which places individuals at risk of infection with pathogens including hepatitis viruses and human immunodeficiency virus (HIV). This notice serves as a reminder that insulin pens must **never** be used on more than one person.

Background

Insulin pens are pen-shaped injector devices that contain a reservoir for insulin or an insulin cartridge. These devices are designed to permit self-injection and are intended for single-person use. In healthcare settings, these devices are often used by healthcare personnel to administer insulin to patients. Insulin pens are designed to be used multiple times, for a single person, using a new needle for each injection. Insulin pens must **never** be used for more than one person. Regurgitation of blood into the insulin cartridge can occur after injection [1] creating a risk of bloodborne pathogen transmission if the pen is used for more than one person, even when the needle is changed.



In 2009, in response to reports of improper use of insulin pens in hospitals, the Food and Drug Administration (FDA) issued an alert for healthcare professionals reminding them that insulin pens are meant for use on a single patient only and are not to be shared between patients [2]. In spite of this alert, there have been continuing reports of patients placed at risk through inappropriate reuse and sharing of insulin pens, including an incident in 2011 that required notification of more than 2,000 potentially exposed patients [3]. These events indicate that some healthcare personnel do not adhere to safe practices and may be unaware of the risks these unsafe practices pose to patients.

Recommendations

Anyone using insulin pens should review the following recommendations to ensure that they are not placing persons in their care at risk for infection.

- Insulin pens containing multiple doses of insulin are meant for use on a single person only, and should **never** be used for more than one person, even when the needle is changed.
- Insulin pens should be clearly labeled with the person's name or other identifying information to ensure that the correct pen is used **only** on the correct individual.
- Hospitals and other facilities should review their policies and educate their staff regarding safe use of insulin pens and similar devices.
- If reuse is identified, exposed persons should be promptly notified and offered appropriate follow-up including bloodborne pathogen testing.

These recommendations apply to any setting where insulin pens are used, including assisted living or residential care facilities, skilled nursing facilities, clinics, health fairs, shelters, detention facilities, senior centers, schools, and camps as well as licensed healthcare facilities. Protection from infections, including bloodborne pathogens, is a basic expectation anywhere healthcare is provided. Use of insulin pens for more than one person, like other forms of syringe reuse [4], imposes unacceptable risks and should be considered a 'never event'.

See additional information on [assuring safe care during blood glucose monitoring and insulin administration](#).

References

1. Sonoki K, Yoshinari M, Iwase M, Tashiro K, Iino K, Wakisaka M, Fujishima M. Regurgitation of blood into insulin cartridges in the pen-like injectors. *Diabetes Care*. 2001;24(3):603-4.
2. [Information for healthcare professionals: risk of transmission of blood-borne pathogens from shared use of insulin pens \(2009\)](#). U.S. Food and Drug Administration Postmarket Drug Safety Information for Patients and Providers.
3. [Important Patient Safety Notification \(2011\)](#). Dean Clinic.
4. [Centers for Disease Control and Prevention \(CDC\) and the Safe Injection Practices Coalition \(SIPC\)](#)

