

**Table 1C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Demographic Indicators** <sup>(2)</sup>

Total Records	Number	%	Source of Data	Number	%
Complete Records	8,775	62.0	WIC	14,142	100.0
Prenatal Records	2,757	19.5	Maternal Health	0	0.0
Postpartum Records	2,610	18.5	Other	0	0.0
<b>Total Records Accepted</b>	<b>14,142</b>		<b>Total</b>	<b>14,142</b>	<b>100.0</b>
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%
White, Not Hispanic	4,457	32.0
Black, Not Hispanic	3,434	24.7
Hispanic	5,445	39.1
American Indian/Alaskan Native	22	0.2
Asian/Pacific Islander	219	1.6
All Other	341	2.5
<b>Total</b>	<b>13,918</b>	<b>100.0</b>
<i>Excluded Unknown: 224</i>		<i>1.6</i>

Age Distribution	Number	%
< 15 Years	70	0.5
15 - 17 Years	951	6.7
18 - 19 Years	1,768	12.5
20 - 29 Years	8,114	57.4
30 - 39 Years	3,010	21.3
>= 40 Years	229	1.6
<b>Total</b>	<b>14,142</b>	<b>100.0</b>
<i>Excluded Unknown: 0</i>		<i>0.0</i>

Education Distribution	Number	%
< High School	0	*
High School	0	*
> High School	0	*
<b>Total</b>	<b>0</b>	<b>100.0</b>
<i>Excluded Unknown: 14,142</i>		<i>100.0</i>

% Poverty Level	Number	%
0 - 50	4,832	40.1
51 - 100	3,059	25.4
101 - 130	1,827	15.2
131 - 150	950	7.9
151 - 185	1,221	10.1
186 - 200	60	0.5
> 200	99	0.8
Adjunctive Eligibility	0	0.0
<b>Total</b>	<b>12,048</b>	<b>100.0</b>
<i>Excluded Unknown: 2,094</i>		<i>14.8</i>

Program Participation <sup>(3)</sup>	Number	%
WIC	11,532	99.1
Food Stamps	11,431	15.4
Medicaid	11,431	68.1
TANF	11,431	11.0
<i>Excluded Unknown:</i>		
<i>WIC:</i>	0	0.0
<i>Food Stamps:</i>	101	0.9
<i>Medicaid:</i>	101	0.9
<i>TANF:</i>	101	0.9

Migrant Status	Number	%
Yes	61	0.4
No	13,566	99.6
<b>Total</b>	<b>13,627</b>	<b>100.0</b>
<i>Excluded Unknown: 515</i>		<i>3.6</i>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Prenatal program participation.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 2C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	13,788	11.7	658,420	12.7
Normal Weight	13,788	47.4	658,420	45.2
Overweight	13,788	14.5	658,420	14.4
Obese	13,788	26.4	658,420	27.8
Maternal Weight Gain (4)				
Less Than Ideal	10,795	24.8	558,067	25.6
Ideal	10,795	29.2	558,067	29.6
Greater Than Ideal	10,795	46.0	558,067	44.8
Anemia (Low Hb/Hct) (5)				
1st Trimester	2,028	4.8	196,351	6.5
2nd Trimester	5,160	4.3	186,011	10.3
3rd Trimester	2,919	26.6	100,071	30.6
Postpartum	8,059	57.6	241,000	28.1
Parity (6)				
0	14,142	45.6	640,776	46.5
1	14,142	28.3	640,776	27.3
2	14,142	15.5	640,776	15.2
≥ 3	14,142	10.6	640,776	11.0
Interpregnancy Interval				
< 6 Months	8,155	11.4	378,012	13.0
6 - <18 Months	8,155	21.5	378,012	27.6
≥18 Months	8,155	67.1	378,012	59.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI < 19.8), normal weight (BMI 19.9 to 26.0), overweight (BMI >26.0 to 29.0), obese (BMI >29.0).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	11,377	84.9	437,601	75.9
No Care	11,377	4.4	437,601	9.6
WIC Enrollment				
1st Trimester	13,888	17.8	666,694	27.4
2nd Trimester	13,888	40.5	666,694	29.4
3rd Trimester	13,888	23.4	666,694	23.4
Postpartum (3)	13,888	18.3	666,694	19.9
Smoking				
3 Months Prior to Pregnancy	14,099	26.0	591,104	30.8
During Pregnancy	11,417	14.2	528,742	19.2
Last 3 Months of Pregnancy	11,385	15.6	423,755	20.3
Postpartum	10,921	16.3	467,521	20.8
Smoking Changes (4)				
Quit Smoking by First Prenatal Visit	3,082	48.6	165,120	41.5
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	2,255	42.0	107,528	31.1
Smoking in Household				
During Pregnancy	11,431	20.8	305,692	27.7
Postpartum	10,929	11.8	340,900	24.0
Drinking				
3 Months Prior to Pregnancy	12,673	4.7	535,600	8.8
Last 3 Months of Pregnancy	11,012	0.3	414,835	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	9,747	0.9	572,813	1.1
Low	9,747	6.5	572,813	6.9
Normal	9,747	85.0	572,813	84.5
High	9,747	7.6	572,813	7.5
Preterm (4)	10,668	9.0	559,297	11.2
Full term Low Birth Weight (5)	8,767	3.7	478,431	4.0
Multiple Births	11,075	0.9	561,253	1.7
Breastfeeding				
Ever Breastfed	10,791	42.1	583,994	55.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.

**Table 3C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Comparison of Racial and Ethnic Distribution by Contributor to Nation**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>							
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
<b>Connecticut</b>	<b>14,142</b>	<b>31.5</b>	<b>24.3</b>	<b>38.5</b>	<b>0.2</b>	<b>1.5</b>	<b>4.0</b>
<b>Nation (Prior Year)</b>	<b>727,815</b>	<b>49.8</b>	<b>24.9</b>	<b>20.1</b>	<b>1.2</b>	<b>2.8</b>	<b>1.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Age Distribution by Contributor to Nation**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
<b>Connecticut</b>	<b>14,142</b>	<b>0.5</b>	<b>6.7</b>	<b>12.5</b>	<b>57.4</b>	<b>21.3</b>	<b>1.6</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>727,815</b>	<b>0.5</b>	<b>7.3</b>	<b>14.1</b>	<b>59.6</b>	<b>17.3</b>	<b>1.2</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Comparison of Maternal Health Indicators by Contributor to Nation**

<b>Comparison of Maternal Health Indicators</b> <sup>(2)</sup>										
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
<b>Connecticut</b>	<b>13,788</b>	<b>11.7</b>	<b>40.9</b>	<b>10,795</b>	<b>24.8</b>	<b>46.0</b>	<b>2,919</b>	<b>26.6</b>	<b>8,059</b>	<b>57.6</b>
<b>Nation (Prior Year)</b>	<b>658,420</b>	<b>12.7</b>	<b>42.2</b>	<b>558,067</b>	<b>25.6</b>	<b>44.8</b>	<b>100,071</b>	<b>30.6</b>	<b>241,000</b>	<b>28.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Maternal Behavioral Indicators by Contributor to Nation**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2)</sup>					
	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
<b>Connecticut</b>	<b>11,377</b>	<b>84.9</b>	<b>4.4</b>	<b>13,888</b>	<b>17.8</b>
<b>Nation (Prior Year)</b>	<b>437,601</b>	<b>75.9</b>	<b>9.6</b>	<b>666,694</b>	<b>27.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Comparison of Smoking Indicators by Contributor to Nation**

<b>Comparison of Smoking Indicators</b> (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
<b>Connecticut</b>	<b>14,099</b>	<b>26.0</b>	<b>11,385</b>	<b>15.6</b>	<b>2,255</b>	<b>42.0</b>	<b>11,431</b>	<b>20.8</b>	<b>10,929</b>	<b>11.8</b>
<b>Nation (Prior Year)</b>	<b>591,104</b>	<b>30.8</b>	<b>423,755</b>	<b>20.3</b>	<b>107,528</b>	<b>31.1</b>	<b>305,692</b>	<b>27.7</b>	<b>340,900</b>	<b>24.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Infant Health Indicators by Contributor to Nation**

<b>Comparison of Infant Health Indicators</b> <sup>(2)</sup>									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
<b>Connecticut</b>	<b>9,747</b>	<b>7.4</b>	<b>7.6</b>	<b>10,668</b>	<b>9.0</b>	<b>8,767</b>	<b>3.7</b>	<b>10,791</b>	<b>42.1</b>
<b>Nation (Prior Year)</b>	<b>572,813</b>	<b>8.0</b>	<b>7.5</b>	<b>559,297</b>	<b>11.2</b>	<b>478,431</b>	<b>4.0</b>	<b>583,994</b>	<b>55.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 9C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Maternal Health Indicators by Race/Ethnicity, Age, or Education**

<b>Maternal Health Indicators by Race/Ethnicity, Age, or Education</b> (2)										
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under- weight	Over- weight		< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number
<b>Race/Ethnicity</b>										
White, Not Hispanic	4,361	13.4	41.2	3,585	20.5	52.5	908	21.5	2,336	54.0
Black, Not Hispanic	3,377	9.4	47.9	2,490	26.5	45.7	855	34.4	2,041	63.4
Hispanic	5,315	10.9	38.0	4,091	26.5	42.1	994	25.7	3,210	57.1
American Indian/Alaskan Native	21	*	*	15	*	*	5	*	10	*
Asian/Pacific Islander	214	31.8	15.9	186	37.1	26.3	56	*	127	52.8
All Other/Unknown	500	13.0	31.8	428	28.5	40.2	101	22.8	335	53.1
<b>Total</b>	<b>13,788</b>	<b>11.7</b>	<b>40.9</b>	<b>10,795</b>	<b>24.8</b>	<b>46.0</b>	<b>2,919</b>	<b>26.6</b>	<b>8,059</b>	<b>57.6</b>
<b>Age</b>										
< 15 Years	70	*	*	46	*	*	21	*	31	*
15 - 17 Years	927	17.6	25.1	684	22.8	48.4	184	28.8	511	59.7
18 - 19 Years	1,724	17.8	28.8	1,256	20.5	52.1	323	28.8	897	62.3
20 - 29 Years	7,903	11.5	41.4	6,156	24.6	46.8	1,737	26.7	4,580	57.6
30 - 39 Years	2,940	6.9	50.7	2,458	27.9	41.0	615	24.6	1,886	55.2
>= 40 Years	224	4.9	60.7	195	29.7	38.5	39	*	154	56.5
Unknown	0	*	*	0	*	*	0	*	0	*
<b>Total</b>	<b>13,788</b>	<b>11.7</b>	<b>40.9</b>	<b>10,795</b>	<b>24.8</b>	<b>46.0</b>	<b>2,919</b>	<b>26.6</b>	<b>8,059</b>	<b>57.6</b>
<b>Education</b>										
< High School	0	*	*	0	*	*	0	*	0	*
High School	0	*	*	0	*	*	0	*	0	*
> High School	0	*	*	0	*	*	0	*	0	*
Unknown	13,788	11.7	40.9	10,795	24.8	46.0	2,919	26.6	8,059	57.6
<b>Total</b>	<b>13,788</b>	<b>11.7</b>	<b>40.9</b>	<b>10,795</b>	<b>24.8</b>	<b>46.0</b>	<b>2,919</b>	<b>26.6</b>	<b>8,059</b>	<b>57.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 10C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education**

<b>Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education</b> (2)					
	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
<b>Race/Ethnicity</b>					
White, Not Hispanic	3,732	87.6	4.6	4,373	17.9
Black, Not Hispanic	2,604	82.6	4.2	3,376	15.8
Hispanic	4,336	84.7	3.7	5,359	19.7
American Indian/Alaskan Native	16	*	*	21	*
Asian/Pacific Islander	192	80.2	5.7	214	8.9
All Other/Unknown	497	80.1	8.7	545	13.9
<b>Total</b>	<b>11,377</b>	<b>84.9</b>	<b>4.4</b>	<b>13,888</b>	<b>17.8</b>
<b>Age</b>					
< 15 Years	47	*	*	70	*
15 - 17 Years	728	76.9	3.7	937	17.9
18 - 19 Years	1,342	81.4	4.0	1,740	20.7
20 - 29 Years	6,491	86.6	4.1	7,963	17.4
30 - 39 Years	2,567	85.2	5.5	2,952	16.8
>= 40 Years	202	83.2	5.0	226	22.1
Unknown	0	*	*	0	*
<b>Total</b>	<b>11,377</b>	<b>84.9</b>	<b>4.4</b>	<b>13,888</b>	<b>17.8</b>
<b>Education</b>					
< High School	0	*	*	0	*
High School	0	*	*	0	*
> High School	0	*	*	0	*
Unknown	11,377	84.9	4.4	13,888	17.8
<b>Total</b>	<b>11,377</b>	<b>84.9</b>	<b>4.4</b>	<b>13,888</b>	<b>17.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 11C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Smoking Indicators by Race/Ethnicity, Age, or Education**

Smoking Indicators by Race/Ethnicity, Age, or Education <sup>(2)</sup>										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
<b>Race/Ethnicity</b>										
White, Not Hispanic	4,447	45.2	3,734	25.9	1,243	37.7	3,352	32.7	3,637	19.1
Black, Not Hispanic	3,406	20.8	2,606	13.0	430	40.9	2,894	19.5	2,517	9.2
Hispanic	5,441	16.1	4,339	9.6	547	50.8	4,680	13.7	4,127	7.6
American Indian/Alaskan Native	22	*	16	*	0	*	15	*	15	*
Asian/Pacific Islander	219	7.3	193	2.6	12	*	150	18.0	190	4.7
All Other/Unknown	564	10.5	497	8.9	23	*	340	14.4	443	8.1
<b>Total</b>	<b>14,099</b>	<b>26.0</b>	<b>11,385</b>	<b>15.6</b>	<b>2,255</b>	<b>42.0</b>	<b>11,431</b>	<b>20.8</b>	<b>10,929</b>	<b>11.8</b>
<b>Age</b>										
< 15 Years	70	*	47	*	4	*	66	*	46	*
15 - 17 Years	945	23.0	728	12.2	140	51.4	842	33.4	691	21.3
18 - 19 Years	1,757	33.4	1,344	18.6	364	49.2	1,517	31.0	1,262	17.8
20 - 29 Years	8,094	26.9	6,493	15.9	1,346	44.0	6,533	19.5	6,233	11.0
30 - 39 Years	3,005	21.4	2,570	14.6	374	26.2	2,308	14.1	2,498	8.2
>= 40 Years	228	18.4	203	15.3	27	*	165	11.5	199	9.5
Unknown	0	*	0	*	0	*	0	*	0	*
<b>Total</b>	<b>14,099</b>	<b>26.0</b>	<b>11,385</b>	<b>15.6</b>	<b>2,255</b>	<b>42.0</b>	<b>11,431</b>	<b>20.8</b>	<b>10,929</b>	<b>11.8</b>
<b>Education</b>										
< High School	0	*	0	*	0	*	0	*	0	*
High School	0	*	0	*	0	*	0	*	0	*
> High School	0	*	0	*	0	*	0	*	0	*
Unknown	14,099	26.0	11,385	15.6	2,255	42.0	11,431	20.8	10,929	11.8
<b>Total</b>	<b>14,099</b>	<b>26.0</b>	<b>11,385</b>	<b>15.6</b>	<b>2,255</b>	<b>42.0</b>	<b>11,431</b>	<b>20.8</b>	<b>10,929</b>	<b>11.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 12C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Infant Health Indicators by Race/Ethnicity, Age, or Education**

<b>Infant Health Indicators by Race/Ethnicity, Age, or Education</b> (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)						
	Number	%	%	Number	%	Number	%	Number	%
<b>Race/Ethnicity</b>									
White, Not Hispanic	3,236	6.2	9.5	3,507	9.1	2,899	2.9	3,574	32.8
Black, Not Hispanic	2,177	9.9	6.0	2,435	10.3	1,931	4.8	2,465	40.2
Hispanic	3,776	7.0	7.2	4,080	8.6	3,426	3.8	4,097	48.9
American Indian/Alaskan Native	11	*	*	12	*	9	*	12	*
Asian/Pacific Islander	170	9.4	0.6	183	7.7	155	3.9	188	47.9
All Other/Unknown	377	5.3	8.2	451	7.1	347	2.6	455	62.0
<b>Total</b>	<b>9,747</b>	<b>7.4</b>	<b>7.6</b>	<b>10,668</b>	<b>9.0</b>	<b>8,767</b>	<b>3.7</b>	<b>10,791</b>	<b>42.1</b>
<b>Age</b>									
< 15 Years	45	*	*	46	*	41	*	46	*
15 - 17 Years	637	6.3	4.2	682	9.8	571	3.5	687	27.7
18 - 19 Years	1,188	7.2	5.7	1,252	8.1	1,074	3.7	1,265	32.6
20 - 29 Years	5,606	6.0	7.9	6,094	8.3	5,077	3.7	6,172	41.8
30 - 39 Years	2,108	7.0	9.1	2,401	10.6	1,870	3.6	2,429	51.1
>= 40 Years	163	10.4	7.4	193	16.1	134	7.5	192	53.6
Unknown	0	*	*	0	*	0	*	0	*
<b>Total</b>	<b>9,747</b>	<b>7.4</b>	<b>7.6</b>	<b>10,668</b>	<b>9.0</b>	<b>8,767</b>	<b>3.7</b>	<b>10,791</b>	<b>42.1</b>
<b>Education</b>									
< High School	0	*	*	0	*	0	*	0	*
High School	0	*	*	0	*	0	*	0	*
> High School	0	*	*	0	*	0	*	0	*
Unknown	9,747	6.5	7.6	10,668	9.0	8,767	3.7	10,791	42.1
<b>Total</b>	<b>9,747</b>	<b>7.4</b>	<b>7.6</b>	<b>10,668</b>	<b>9.0</b>	<b>8,767</b>	<b>3.7</b>	<b>10,791</b>	<b>42.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 13C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Maternal Weight Gain and Birth Outcomes by Select Health Indicators**

<b>Maternal Weight Gain and Birth Outcomes by Select Health Indicators</b> (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)	Full Term LBW (7)		
		Number	%	%	Low (4)	High (5)				
	Number	%	%	Number	%	%	Number	%	Number	%
<b>Prepregnancy BMI</b>										
Underweight	1,242	26.8	33.4	1,102	9.8	2.9	1,208	9.4	988	5.3
Normal Weight	5,097	25.6	43.0	4,514	7.3	6.8	4,932	8.6	4,080	3.9
Overweight	1,588	15.8	61.5	1,395	5.8	9.7	1,538	8.3	1,267	2.9
Obese	2,865	27.5	48.2	2,577	7.0	9.9	2,762	9.8	2,293	2.8
<b>Weight Gain</b>										
< Ideal	-	-	-	2,316	12.1	3.8	2,534	12.5	1,990	5.8
Ideal	-	-	-	2,806	7.2	5.9	3,062	9.5	2,518	3.5
> Ideal	-	-	-	4,430	4.8	10.7	4,803	6.8	4,088	2.7
<b>Smoking During Pregnancy</b>										
Smokers	1,136	26.2	46.3	1,078	11.6	4.8	1,102	11.9	926	5.8
Non-Smokers	7,442	24.4	46.0	7,083	6.5	7.9	7,277	8.1	6,445	3.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 14C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Racial and Ethnic Distribution**

<b>Summary of Trends in Racial and Ethnic Distribution</b> <sup>(2)</sup>							
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
2003	14,142	31.5	24.3	38.5	0.2	1.5	4.0
2002	0	*	*	*	*	*	*
2001	10,360	34.4	23.8	35.9	0.3	1.6	4.1
2000	0	*	*	*	*	*	*
1999	0	*	*	*	*	*	*
1998	0	*	*	*	*	*	*
1997	0	*	*	*	*	*	*
1996	0	*	*	*	*	*	*
1995	0	*	*	*	*	*	*
1994	0	*	*	*	*	*	*
1993	0	*	*	*	*	*	*
1992	0	*	*	*	*	*	*
1991	0	*	*	*	*	*	*
1990	0	*	*	*	*	*	*
1989	409	48.4	32.0	19.1	0.5	0.0	0.0
1988	812	46.7	30.4	22.0	0.5	0.4	0.0
1987	555	49.2	31.5	18.9	0.2	0.2	0.0
1986	331	58.0	29.0	12.7	0.3	0.0	0.0
1985	545	51.0	31.0	16.5	1.3	0.2	0.0
1984	534	58.1	29.0	11.8	0.9	0.2	0.0
1983	430	58.8	26.7	13.7	0.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 15C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Age Distribution**

<b>Summary of Trends in Age Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2003	14,142	0.5	6.7	12.5	57.4	21.3	1.6	0.0
2002	0	*	*	*	*	*	*	*
2001	10,360	0.5	7.2	12.1	54.9	23.4	1.9	0.0
2000	0	*	*	*	*	*	*	*
1999	0	*	*	*	*	*	*	*
1998	0	*	*	*	*	*	*	*
1997	0	*	*	*	*	*	*	*
1996	0	*	*	*	*	*	*	*
1995	0	*	*	*	*	*	*	*
1994	0	*	*	*	*	*	*	*
1993	0	*	*	*	*	*	*	*
1992	0	*	*	*	*	*	*	*
1991	0	*	*	*	*	*	*	*
1990	0	*	*	*	*	*	*	*
1989	409	0.5	10.0	14.4	57.7	17.1	0.2	0.0
1988	812	0.4	12.3	14.0	56.7	15.5	1.1	0.0
1987	555	0.7	8.1	12.6	60.0	17.8	0.7	0.0
1986	331	0.9	12.1	16.3	58.3	12.1	0.3	0.0
1985	545	1.5	12.8	17.6	51.6	15.6	0.9	0.0
1984	534	0.6	12.0	17.6	56.4	12.4	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 15C**  
**1983 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Age Distribution**

<b>Summary of Trends in Age Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1983	430	0.5	12.8	16.3	59.3	10.9	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 16C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Health Indicators**

Summary of Trends in Maternal Health Indicators <sup>(2)</sup>										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2003	13,788	11.7	40.9	10,795	24.8	46.0	2,919	26.6	8,059	57.6
2002	0	*	*	0	*	*	0	*	0	*
2001	10,215	11.8	41.0	10,171	26.7	43.3	1,987	29.9	7,737	57.9
2000	0	*	*	0	*	*	0	*	0	*
1999	0	*	*	0	*	*	0	*	0	*
1998	0	*	*	0	*	*	0	*	0	*
1997	0	*	*	0	*	*	0	*	0	*
1996	0	*	*	0	*	*	0	*	0	*
1995	0	*	*	0	*	*	0	*	0	*
1994	0	*	*	0	*	*	0	*	0	*
1993	0	*	*	0	*	*	0	*	0	*
1992	0	*	*	0	*	*	0	*	0	*
1991	0	*	*	0	*	*	0	*	0	*
1990	0	*	*	0	*	*	0	*	0	*
1989	408	20.1	23.5	0	*	*	99	*	136	34.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 16C**  
**1988 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Health Indicators**

Summary of Trends in Maternal Health Indicators <sup>(2)</sup>										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1988	789	18.0	28.3	0	*	*	132	22.7	152	39.5
1987	549	19.7	26.8	0	*	*	134	34.3	126	24.6
1986	325	19.4	24.9	0	*	*	67	*	107	23.4
1985	541	22.0	22.4	0	*	*	118	19.5	214	20.1
1984	533	22.7	18.4	0	*	*	126	25.4	255	21.2
1983	426	20.7	21.4	0	*	*	96	*	217	26.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 17C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Behavioral Indicators**

<b>Summary of Trends in Maternal Behavioral Indicators</b> <sup>(2)</sup>					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2003	11,377	84.9	4.4	13,888	17.8
2002	0	*	*	0	*
2001	10,331	81.8	6.8	10,160	17.9
2000	0	*	*	0	*
1999	0	*	*	0	*
1998	0	*	*	0	*
1997	0	*	*	0	*
1996	0	*	*	0	*
1995	0	*	*	0	*
1994	0	*	*	0	*
1993	0	*	*	0	*
1992	0	*	*	0	*
1991	0	*	*	0	*
1990	0	*	*	0	*
1989	0	*	*	0	*
1988	0	*	*	0	*
1987	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 17C**  
**1986 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Behavioral Indicators**

<b>Summary of Trends in Maternal Behavioral Indicators</b> <sup>(2)</sup>					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
1986	0	*	*	0	*
1985	0	*	*	0	*
1984	0	*	*	0	*
1983	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 18C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators**

<b>Summary of Trends in Smoking Indicators</b> (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2003	14,099	26.0	11,385	15.6	2,255	42.0	11,431	20.8	10,929	11.8
2002	0	*	0	*	0	*	0	*	0	*
2001	10,320	27.1	10,346	15.3	2,308	37.9	8,079	24.5	10,271	14.8
2000	0	*	0	*	0	*	0	*	0	*
1999	0	*	0	*	0	*	0	*	0	*
1998	0	*	0	*	0	*	0	*	0	*
1997	0	*	0	*	0	*	0	*	0	*
1996	0	*	0	*	0	*	0	*	0	*
1995	0	*	0	*	0	*	0	*	0	*
1994	0	*	0	*	0	*	0	*	0	*
1993	0	*	0	*	0	*	0	*	0	*
1992	0	*	0	*	0	*	0	*	0	*
1991	0	*	0	*	0	*	0	*	0	*
1990	0	*	0	*	0	*	0	*	0	*
1989	0	*	0	*	0	*	0	*	0	*
1988	0	*	0	*	0	*	0	*	0	*
1987	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 18C**  
**1986 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators**

<b>Summary of Trends in Smoking Indicators</b> (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
1986	0	*	0	*	0	*	0	*	0	*
1985	0	*	0	*	0	*	0	*	0	*
1984	0	*	0	*	0	*	0	*	0	*
1983	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 19C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Infant Health Indicators**

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2003	9,747	7.4	7.6	10,668	9.0	8,767	3.7	10,791	42.1
2002	0	*	*	0	*	0	*	0	*
2001	9,016	7.1	8.6	9,985	8.8	8,113	3.5	10,093	40.0
2000	0	*	*	0	*	0	*	0	*
1999	0	*	*	0	*	0	*	0	*
1998	0	*	*	0	*	0	*	0	*
1997	0	*	*	0	*	0	*	0	*
1996	0	*	*	0	*	0	*	0	*
1995	0	*	*	0	*	0	*	0	*
1994	0	*	*	0	*	0	*	0	*
1993	0	*	*	0	*	0	*	0	*
1992	0	*	*	0	*	0	*	0	*
1991	0	*	*	0	*	0	*	0	*
1990	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 19C**  
**1989 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Infant Health Indicators**

<b>Summary of Trends in Infant Health Indicators</b> (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1989	409	7.6	7.1	400	0.5	398	7.5	100	100.0
1988	411	5.8	11.4	409	0.7	406	5.7	95	*
1987	354	6.2	9.6	350	0.9	347	6.1	63	*
1986	324	7.4	11.1	322	1.6	317	7.3	65	*
1985	525	6.1	9.3	522	0.8	518	6.0	99	*
1984	500	9.2	9.4	495	2.4	483	8.9	149	100.0
1983	424	7.3	6.4	422	4.0	405	6.7	123	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2003	4,361	13.4	41.2	3,585	20.5	52.5	908	21.5	2,336	54.0
	2002	0	*	*	0	*	*	0	*	0	*
	2001	3,527	14.1	41.1	3,510	23.0	47.2	673	24.8	2,373	55.0
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*
Black, Not Hispanic	2003	3,377	9.4	47.9	2,490	26.5	45.7	855	34.4	2,041	63.4
	2002	0	*	*	0	*	*	0	*	0	*
	2001	2,430	8.7	50.7	2,413	28.9	43.3	565	38.2	2,019	63.6
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2003	5,315	10.9	38.0	4,091	26.5	42.1	994	25.7	3,210	57.1
	2002	0	*	*	0	*	*	0	*	0	*
	2001	3,673	10.5	37.2	3,663	27.7	40.9	638	29.3	2,900	57.4
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*
American Indian/Alaskan Native	2003	21	*	*	15	*	*	5	*	10	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	26	*	*	26	*	*	7	*	21	*
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2003 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2003	214	31.8	15.9	186	37.1	26.3	56	*	127	52.8
	2002	0	*	*	0	*	*	0	*	0	*
	2001	166	28.9	15.1	166	30.7	34.3	31	*	119	55.5
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*
All Other/Unknown	2003	500	13.0	31.8	428	28.5	40.2	101	22.8	335	53.1
	2002	0	*	*	0	*	*	0	*	0	*
	2001	393	15.3	27.2	393	34.4	36.6	73	*	305	49.5
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2003 Pregnancy Nutrition Surveillance (1)**  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity (2)</b>											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Total All Race/Ethnic Groups	2003	13,788	11.7	40.9	10,795	24.8	46.0	2,919	26.6	8,059	57.6
	2002	0	*	*	0	*	*	0	*	0	*
	2001	10,215	11.8	41.0	10,171	26.7	43.3	1,987	29.9	7,737	57.9
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
	1995	0	*	*	0	*	*	0	*	0	*
	1994	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2003	3,732	87.6	4.6	4,373	17.9
	2002	0	*	*	0	*
	2001	3,551	86.3	5.7	3,488	18.1
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*
Black, Not Hispanic	2003	2,604	82.6	4.2	3,376	15.8
	2002	0	*	*	0	*
	2001	2,456	79.2	7.0	2,415	16.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2003	4,336	84.7	3.7	5,359	19.7
	2002	0	*	*	0	*
	2001	3,708	80.3	7.0	3,652	19.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*
American Indian/Alaskan Native	2003	16	*	*	21	*
	2002	0	*	*	0	*
	2001	26	*	*	26	*
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2003	192	80.2	5.7	214	8.9
	2002	0	*	*	0	*
	2001	168	78.0	10.7	165	11.5
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*
All Other/Unknown	2003	497	80.1	8.7	545	13.9
	2002	0	*	*	0	*
	2001	422	74.4	12.3	414	11.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Total All Race/Ethnic Groups	2003	11,377	84.9	4.4	13,888	17.8
	2002	0	*	*	0	*
	2001	10,331	81.8	6.8	10,160	17.9
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
	1995	0	*	*	0	*
	1994	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2003 Pregnancy Nutrition Surveillance** (1)

**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2003	4,447	45.2	3,734	25.9	1,243	37.7	3,352	32.7	3,637	19.1
	2002	0	*	0	*	0	*	0	*	0	*
	2001	3,550	46.8	3,556	28.3	1,306	34.3	2,569	38.3	3,540	23.2
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*
Black, Not Hispanic	2003	3,406	20.8	2,606	13.0	430	40.9	2,894	19.5	2,517	9.2
	2002	0	*	0	*	0	*	0	*	0	*
	2001	2,447	21.8	2,462	12.1	481	37.4	2,005	20.5	2,450	11.6
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2003 Pregnancy Nutrition Surveillance** (1)

**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2003	5,441	16.1	4,339	9.6	547	50.8	4,680	13.7	4,127	7.6
	2002	0	*	0	*	0	*	0	*	0	*
	2001	3,708	14.7	3,712	6.8	486	47.3	3,108	16.7	3,688	9.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*
American Indian/Alaskan Native	2003	22	*	16	*	0	*	15	*	15	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	26	*	26	*	2	*	20	*	26	*
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2003 Pregnancy Nutrition Surveillance** (1)

**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

Summary of Trends in Smoking Indicators by Race/Ethnicity (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2003	219	7.3	193	2.6	12	*	150	18.0	190	4.7
	2002	0	*	0	*	0	*	0	*	0	*
	2001	168	6.5	168	2.4	9	*	109	17.4	168	12.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*
All Other/Unknown	2003	564	10.5	497	8.9	23	*	340	14.4	443	8.1
	2002	0	*	0	*	0	*	0	*	0	*
	2001	421	10.0	422	6.4	24	*	268	15.7	399	8.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2003 Pregnancy Nutrition Surveillance** (1)

**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Total All Race/Ethnic Groups	2003	14,099	26.0	11,385	15.6	2,255	42.0	11,431	20.8	10,929	11.8
	2002	0	*	0	*	0	*	0	*	0	*
	2001	10,320	27.1	10,346	15.3	2,308	37.9	8,079	24.5	10,271	14.8
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
	1995	0	*	0	*	0	*	0	*	0	*
	1994	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2003	3,236	6.2	9.5	3,507	9.1	2,899	2.9	3,574	32.8
	2002	0	*	*	0	*	0	*	0	*
	2001	3,150	6.0	10.2	3,419	8.2	2,845	2.8	3,470	31.5
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*
Black, Not Hispanic	2003	2,177	9.9	6.0	2,435	10.3	1,931	4.8	2,465	40.2
	2002	0	*	*	0	*	0	*	0	*
	2001	2,103	9.9	6.3	2,365	9.7	1,867	5.2	2,387	35.7
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
Hispanic	2003	3,776	7.0	7.2	4,080	8.6	3,426	3.8	4,097	48.9
	2002	0	*	*	0	*	0	*	0	*
	2001	3,251	6.2	8.9	3,603	8.9	2,941	3.1	3,634	48.0
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*
American Indian/Alaskan Native	2003	11	*	*	12	*	9	*	12	*
	2002	0	*	*	0	*	0	*	0	*
	2001	22	*	*	26	*	18	*	25	*
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
Asian/Pacific Islander	2003	170	9.4	0.6	183	7.7	155	3.9	188	47.9
	2002	0	*	*	0	*	0	*	0	*
	2001	140	7.1	5.7	163	9.2	125	2.4	164	42.7
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*
All Other/Unknown	2003	377	5.3	8.2	451	7.1	347	2.6	455	62.0
	2002	0	*	*	0	*	0	*	0	*
	2001	350	7.1	7.4	409	8.1	317	3.2	413	62.2
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2003 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)	Number	%	Number	%	Number	%
		Number	%	%						
Total All Race/Ethnic Groups	2003	9,747	7.4	7.6	10,668	9.0	8,767	3.7	10,791	42.1
	2002	0	*	*	0	*	0	*	0	*
	2001	9,016	7.1	8.6	9,985	8.8	8,113	3.5	10,093	40.0
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
	1995	0	*	*	0	*	0	*	0	*
	1994	0	*	*	0	*	0	*	0	*

- (1) Reporting period is January 1 through December 31.
- (2) Excludes data with errors.
- (3) <2500 g.
- (4) >4000 g.
- (5) <37 weeks gestation.
- (6) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.