

**Table 1C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Demographic Indicators** <sup>(2)</sup>

Total Records	Number	%	Source of Data	Number	%
Complete Records	8,611	81.1	WIC	10,624	100.0
Prenatal Records	482	4.5	Maternal Health	0	0.0
Postpartum Records	1,531	14.4	Other	0	0.0
<b>Total Records Accepted</b>	<b>10,624</b>		<b>Total</b>	<b>10,624</b>	<b>100.0</b>
			<i>Excluded Unknown:</i> 0		0.0

Race/Ethnic Distribution	Number	%	Age Distribution	Number	%	Education Distribution	Number	%
White, Not Hispanic	3,509	33.1	< 15 Years	37	0.3	< High School	0	*
Black, Not Hispanic	2,331	22.0	15 - 17 Years	687	6.5	High School	0	*
Hispanic	4,466	42.1	18 - 19 Years	1,216	11.4	> High School	0	*
American Indian/Alaskan Native	45	0.4	20 - 29 Years	6,114	57.5	<b>Total</b>	<b>0</b>	<b>100.0</b>
Asian/Pacific Islander	227	2.1	30 - 39 Years	2,367	22.3	<i>Excluded Unknown:</i> 10,624		100.0
All Other	18	0.2	>= 40 Years	203	1.9			
<b>Total</b>	<b>10,596</b>	<b>100.0</b>	<b>Total</b>	<b>10,624</b>	<b>100.0</b>			
<i>Excluded Unknown:</i> 28		0.3	<i>Excluded Unknown:</i> 0		0.0			

% Poverty Level	Number	%	Program Participation <sup>(3)</sup>	Number	%	Migrant Status	Number	%
0 - 50	2,460	34.3	WIC	9,093	100.0	Yes	5	0.0
51 - 100	2,065	28.8	Food Stamps	0	*	No	10,256	100.0
101 - 130	1,163	16.2	Medicaid	0	*	<b>Total</b>	<b>10,261</b>	<b>100.0</b>
131 - 150	513	7.2	TANF	0	*	<i>Excluded Unknown:</i> 363		3.4
151 - 185	632	8.8	<i>Excluded Unknown:</i>					
186 - 200	56	0.8	WIC:	0	0.0			
> 200	275	3.8	Food Stamps:	9,093	100.0			
Adjunctive Eligibility	0	0.0	Medicaid:	9,093	100.0			
<b>Total</b>	<b>7,164</b>	<b>100.0</b>	TANF:	9,093	100.0			
<i>Excluded Unknown:</i> 3,460		32.6						

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 2C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	10,536	9.5	784,438	11.7
Normal Weight	10,536	46.2	784,438	44.9
Overweight	10,536	15.7	784,438	14.6
Obese	10,536	28.6	784,438	28.7
Maternal Weight Gain (4)				
Less Than Ideal	10,066	69.7	650,654	25.6
Ideal	10,066	17.4	650,654	30.7
Greater Than Ideal	10,066	12.9	650,654	43.7
Anemia (Low Hb/Hct) (5)				
1st Trimester	1,661	4.8	230,960	6.7
2nd Trimester	3,845	4.6	229,832	10.5
3rd Trimester	1,956	23.2	119,915	30.7
Postpartum	7,271	58.1	254,974	29.4
Parity (6)				
0	6,890	0.0	773,919	47.9
1	6,890	42.3	773,919	25.1
2	6,890	26.8	773,919	14.7
≥ 3	6,890	30.9	773,919	12.4
Interpregnancy Interval				
< 6 Months	6,390	12.9	433,196	13.1
6 - <18 Months	6,390	23.0	433,196	27.2
≥18 Months	6,390	64.1	433,196	59.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI < 19.8), normal weight (BMI 19.9 to 26.0), overweight (BMI >26.0 to 29.0), obese (BMI >29.0).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	10,135	86.7	500,728	76.7
No Care	10,135	2.8	500,728	8.1
WIC Enrollment				
1st Trimester	10,264	21.9	718,544	30.9
2nd Trimester	10,264	42.3	718,544	34.0
3rd Trimester	10,264	21.2	718,544	18.4
Postpartum (3)	10,264	14.6	718,544	16.6
Smoking				
3 Months Prior to Pregnancy	10,605	27.3	712,987	32.6
During Pregnancy	9,081	12.5	627,967	17.5
Last 3 Months of Pregnancy	10,142	3.7	592,858	17.9
Postpartum	10,141	15.0	585,032	18.7
Smoking Changes (4)				
Quit Smoking by First Prenatal Visit	2,492	54.5	202,180	48.7
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	2,358	53.7	156,004	40.8
Smoking in Household				
During Pregnancy	9,093	16.9	368,918	25.6
Postpartum	10,142	9.6	460,583	26.4
Drinking				
3 Months Prior to Pregnancy	1	*	325,168	9.9
Last 3 Months of Pregnancy	0	*	540,106	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 2C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Health Indicators**

Summary of Health Indicators (2)				
	Connecticut Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	10,126	1.1	681,087	1.2
Low	10,126	6.7	681,087	7.0
Normal	10,126	84.6	681,087	84.6
High	10,126	7.6	681,087	7.2
Preterm (4)	9,763	8.9	618,497	12.5
Full term Low Birth Weight (5)	8,888	3.5	557,817	3.9
Multiple Births	10,142	0.8	734,555	1.4
Breastfeeding				
Ever Breastfed	10,137	61.9	698,846	57.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.

**Table 3C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Racial and Ethnic Distribution by Contributor to Nation**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>							
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
<b>Connecticut</b>	<b>10,624</b>	<b>33.0</b>	<b>21.9</b>	<b>42.0</b>	<b>0.4</b>	<b>2.1</b>	<b>0.4</b>
<b>Nation (Prior Year)</b>	<b>856,123</b>	<b>48.0</b>	<b>23.3</b>	<b>21.7</b>	<b>1.4</b>	<b>2.2</b>	<b>3.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Age Distribution by Contributor to Nation**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
<b>Connecticut</b>	<b>10,624</b>	<b>0.3</b>	<b>6.5</b>	<b>11.4</b>	<b>57.5</b>	<b>22.3</b>	<b>1.9</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>856,123</b>	<b>0.4</b>	<b>6.8</b>	<b>13.2</b>	<b>60.3</b>	<b>18.0</b>	<b>1.3</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Maternal Health Indicators by Contributor to Nation**

<b>Comparison of Maternal Health Indicators</b> <sup>(2)</sup>										
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
<b>Connecticut</b>	<b>10,536</b>	<b>9.5</b>	<b>44.2</b>	<b>10,066</b>	<b>69.7</b>	<b>12.9</b>	<b>1,956</b>	<b>23.2</b>	<b>7,271</b>	<b>58.1</b>
<b>Nation (Prior Year)</b>	<b>784,438</b>	<b>11.7</b>	<b>43.3</b>	<b>650,654</b>	<b>25.6</b>	<b>43.7</b>	<b>119,915</b>	<b>30.7</b>	<b>254,974</b>	<b>29.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Maternal Behavioral Indicators by Contributor to Nation**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2)</sup>					
	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
<b>Connecticut</b>	<b>10,135</b>	<b>86.7</b>	<b>2.8</b>	<b>10,264</b>	<b>21.9</b>
<b>Nation (Prior Year)</b>	<b>500,728</b>	<b>76.7</b>	<b>8.1</b>	<b>718,544</b>	<b>30.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Comparison of Smoking Indicators by Contributor to Nation**

<b>Comparison of Smoking Indicators</b> (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
<b>Connecticut</b>	<b>10,605</b>	<b>27.3</b>	<b>10,142</b>	<b>3.7</b>	<b>2,358</b>	<b>53.7</b>	<b>9,093</b>	<b>16.9</b>	<b>10,142</b>	<b>9.6</b>
<b>Nation (Prior Year)</b>	<b>712,987</b>	<b>32.6</b>	<b>592,858</b>	<b>17.9</b>	<b>156,004</b>	<b>40.8</b>	<b>368,918</b>	<b>25.6</b>	<b>460,583</b>	<b>26.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Comparison of Infant Health Indicators by Contributor to Nation**

<b>Comparison of Infant Health Indicators</b> <sup>(2)</sup>									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)	Number	%	Number	%	Number	%
	Number	%	%						
<b>Connecticut</b>	<b>10,126</b>	<b>7.7</b>	<b>7.6</b>	<b>9,763</b>	<b>8.9</b>	<b>8,888</b>	<b>3.5</b>	<b>10,137</b>	<b>61.9</b>
<b>Nation (Prior Year)</b>	<b>681,087</b>	<b>8.2</b>	<b>7.2</b>	<b>618,497</b>	<b>12.5</b>	<b>557,817</b>	<b>3.9</b>	<b>698,846</b>	<b>57.3</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 9C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Maternal Health Indicators by Race/Ethnicity, Age, or Education**

<b>Maternal Health Indicators by Race/Ethnicity, Age, or Education</b> <sup>(2)</sup>										
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
<b>Race/Ethnicity</b>										
White, Not Hispanic	3,466	11.1	44.1	3,351	62.8	18.8	644	17.9	2,102	54.7
Black, Not Hispanic	2,314	7.4	51.4	2,181	70.8	12.1	497	31.8	1,711	62.2
Hispanic	4,442	8.8	41.8	4,228	74.3	8.9	757	21.8	3,254	58.5
American Indian/Alaskan Native	45	*	*	41	*	*	8	*	27	*
Asian/Pacific Islander	223	18.8	25.6	219	79.5	7.3	43	*	147	53.1
All Other/Unknown	46	*	*	46	*	*	7	*	30	*
<b>Total</b>	<b>10,536</b>	<b>9.5</b>	<b>44.2</b>	<b>10,066</b>	<b>69.7</b>	<b>12.9</b>	<b>1,956</b>	<b>23.2</b>	<b>7,271</b>	<b>58.1</b>
<b>Age</b>										
< 15 Years	37	*	*	34	*	*	8	*	31	*
15 - 17 Years	684	15.6	25.7	639	67.4	14.9	109	31.2	482	63.7
18 - 19 Years	1,211	15.4	33.8	1,142	65.4	16.4	185	20.0	807	62.2
20 - 29 Years	6,063	9.2	44.5	5,805	68.6	13.3	1,152	24.1	4,114	57.9
30 - 39 Years	2,341	5.8	54.0	2,253	74.6	10.2	458	20.7	1,689	55.2
>= 40 Years	200	4.0	53.0	193	79.8	5.2	44	*	148	56.1
Unknown	0	*	*	0	*	*	0	*	0	*
<b>Total</b>	<b>10,536</b>	<b>9.5</b>	<b>44.2</b>	<b>10,066</b>	<b>69.7</b>	<b>12.9</b>	<b>1,956</b>	<b>23.2</b>	<b>7,271</b>	<b>58.1</b>
<b>Education</b>										
< High School	0	*	*	0	*	*	0	*	0	*
High School	0	*	*	0	*	*	0	*	0	*
> High School	0	*	*	0	*	*	0	*	0	*
Unknown	10,536	9.5	44.2	10,066	69.7	12.9	1,956	23.2	7,271	58.1
<b>Total</b>	<b>10,536</b>	<b>9.5</b>	<b>44.2</b>	<b>10,066</b>	<b>69.7</b>	<b>12.9</b>	<b>1,956</b>	<b>23.2</b>	<b>7,271</b>	<b>58.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 10C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education**

<b>Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education</b> <sup>(2)</sup>					
	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
<b>Race/Ethnicity</b>					
White, Not Hispanic	3,385	88.5	3.4	3,378	23.0
Black, Not Hispanic	2,190	84.4	2.5	2,270	19.4
Hispanic	4,250	86.8	2.3	4,315	22.7
American Indian/Alaskan Native	41	*	*	42	*
Asian/Pacific Islander	223	78.5	6.3	213	14.1
All Other/Unknown	46	*	*	46	*
<b>Total</b>	<b>10,135</b>	<b>86.7</b>	<b>2.8</b>	<b>10,264</b>	<b>21.9</b>
<b>Age</b>					
< 15 Years	34	*	*	35	*
15 - 17 Years	641	80.5	1.7	661	27.1
18 - 19 Years	1,147	86.1	2.5	1,152	25.8
20 - 29 Years	5,846	87.3	2.6	5,903	21.1
30 - 39 Years	2,271	87.3	3.7	2,310	21.3
>= 40 Years	196	86.2	5.1	203	13.8
Unknown	0	*	*	0	*
<b>Total</b>	<b>10,135</b>	<b>86.7</b>	<b>2.8</b>	<b>10,264</b>	<b>21.9</b>
<b>Education</b>					
< High School	0	*	*	0	*
High School	0	*	*	0	*
> High School	0	*	*	0	*
Unknown	10,135	86.7	2.8	10,264	21.9
<b>Total</b>	<b>10,135</b>	<b>86.7</b>	<b>2.8</b>	<b>10,264</b>	<b>21.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 11C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Smoking Indicators by Race/Ethnicity, Age, or Education**

<b>Smoking Indicators by Race/Ethnicity, Age, or Education</b> <sup>(2)</sup>										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
<b>Race/Ethnicity</b>										
White, Not Hispanic	3,508	45.3	3,390	8.3	1,240	47.3	2,739	26.5	3,390	15.8
Black, Not Hispanic	2,319	23.6	2,191	2.0	475	54.9	2,110	16.5	2,191	7.9
Hispanic	4,460	16.1	4,251	1.1	614	65.3	4,002	10.6	4,251	5.6
American Indian/Alaskan Native	45	*	41	*	8	*	34	*	41	*
Asian/Pacific Islander	227	5.3	223	0.4	11	*	173	12.1	223	9.9
All Other/Unknown	46	*	46	*	10	*	35	*	46	*
<b>Total</b>	<b>10,605</b>	<b>27.3</b>	<b>10,142</b>	<b>3.7</b>	<b>2,358</b>	<b>53.7</b>	<b>9,093</b>	<b>16.9</b>	<b>10,142</b>	<b>9.6</b>
<b>Age</b>										
< 15 Years	37	*	34	*	3	*	36	*	34	*
15 - 17 Years	685	23.9	642	2.2	149	58.4	649	30.7	642	18.1
18 - 19 Years	1,215	33.3	1,147	3.0	343	58.9	1,079	27.0	1,147	15.3
20 - 29 Years	6,099	29.3	5,849	3.4	1,425	56.6	5,229	15.4	5,849	9.0
30 - 39 Years	2,366	21.2	2,274	5.3	412	39.1	1,936	10.8	2,274	6.3
>= 40 Years	203	15.3	196	4.6	26	*	164	12.8	196	6.1
Unknown	0	*	0	*	0	*	0	*	0	*
<b>Total</b>	<b>10,605</b>	<b>27.3</b>	<b>10,142</b>	<b>3.7</b>	<b>2,358</b>	<b>53.7</b>	<b>9,093</b>	<b>16.9</b>	<b>10,142</b>	<b>9.6</b>
<b>Education</b>										
< High School	0	*	0	*	0	*	0	*	0	*
High School	0	*	0	*	0	*	0	*	0	*
> High School	0	*	0	*	0	*	0	*	0	*
Unknown	10,605	27.3	10,142	3.7	2,358	53.7	9,093	16.9	10,142	9.6
<b>Total</b>	<b>10,605</b>	<b>27.3</b>	<b>10,142</b>	<b>3.7</b>	<b>2,358</b>	<b>53.7</b>	<b>9,093</b>	<b>16.9</b>	<b>10,142</b>	<b>9.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 12C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Infant Health Indicators by Race/Ethnicity, Age, or Education**

<b>Infant Health Indicators by Race/Ethnicity, Age, or Education</b> (2)									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)	Number	%	Number	%	Number	%
	Number	%	%						
<b>Race/Ethnicity</b>									
White, Not Hispanic	3,385	7.1	9.1	3,235	9.1	2,940	3.1	3,387	56.3
Black, Not Hispanic	2,185	10.9	6.1	2,137	10.5	1,909	4.8	2,190	54.4
Hispanic	4,246	6.5	7.5	4,100	8.0	3,770	3.0	4,250	69.5
American Indian/Alaskan Native	41	*	*	36	*	33	*	41	*
Asian/Pacific Islander	223	9.0	4.5	209	7.7	193	5.2	223	73.5
All Other/Unknown	46	*	*	46	*	43	*	46	*
<b>Total</b>	<b>10,126</b>	<b>7.7</b>	<b>7.6</b>	<b>9,763</b>	<b>8.9</b>	<b>8,888</b>	<b>3.5</b>	<b>10,137</b>	<b>61.9</b>
<b>Age</b>									
< 15 Years	34	*	*	32	*	31	*	34	*
15 - 17 Years	642	7.5	3.6	618	8.7	564	4.3	642	51.1
18 - 19 Years	1,145	7.9	5.1	1,081	8.5	987	4.5	1,147	59.2
20 - 29 Years	5,839	5.9	7.9	5,622	7.8	5,180	3.0	5,845	62.2
30 - 39 Years	2,271	7.7	9.5	2,214	11.4	1,959	4.0	2,273	65.1
>= 40 Years	195	7.7	7.2	196	14.8	167	3.6	196	71.9
Unknown	0	*	*	0	*	0	*	0	*
<b>Total</b>	<b>10,126</b>	<b>7.7</b>	<b>7.6</b>	<b>9,763</b>	<b>8.9</b>	<b>8,888</b>	<b>3.5</b>	<b>10,137</b>	<b>61.9</b>
<b>Education</b>									
< High School	0	*	*	0	*	0	*	0	*
High School	0	*	*	0	*	0	*	0	*
> High School	0	*	*	0	*	0	*	0	*
Unknown	10,126	6.7	7.6	9,763	8.9	8,888	3.5	10,137	61.9
<b>Total</b>	<b>10,126</b>	<b>7.7</b>	<b>7.6</b>	<b>9,763</b>	<b>8.9</b>	<b>8,888</b>	<b>3.5</b>	<b>10,137</b>	<b>61.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 13C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Maternal Weight Gain and Birth Outcomes by Select Health Indicators**

<b>Maternal Weight Gain and Birth Outcomes by Select Health Indicators</b> (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal	> Ideal		Birthweight			Preterm (6)	Full Term LBW (7)		
		Low (4)	High (5)							
Number	%	%	Number	%	%	Number	%	Number	%	
<b>Prepregnancy BMI</b>										
Underweight	952	76.9	6.2	951	10.5	3.3	913	8.5	834	6.6
Normal Weight	4,645	73.9	10.2	4,637	7.8	5.9	4,505	8.6	4,112	3.4
Overweight	1,571	57.8	20.2	1,569	6.9	9.5	1,502	9.5	1,359	2.8
Obese	2,884	67.1	15.5	2,882	7.0	10.9	2,756	9.1	2,502	3.0
<b>Weight Gain</b>										
< Ideal	-	-	-	7,003	8.2	6.6	6,697	9.3	6,068	3.7
Ideal	-	-	-	1,748	7.0	8.4	1,713	8.6	1,565	2.9
> Ideal	-	-	-	1,300	6.2	12.2	1,278	7.3	1,184	3.1
<b>Smoking During Pregnancy</b>										
Smokers	1,070	65.3	17.7	1,072	10.0	4.6	1,034	10.0	930	4.7
Non-Smokers	7,519	71.8	11.0	7,514	6.9	8.2	7,268	8.2	6,664	3.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 14C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Racial and Ethnic Distribution**

<b>Summary of Trends in Racial and Ethnic Distribution</b> <sup>(2)</sup>							
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
2005	10,624	33.0	21.9	42.0	0.4	2.1	0.4
2004	10,219	33.1	23.1	38.6	0.2	1.8	3.2
2003	14,142	31.5	24.3	38.5	0.2	1.5	4.0
2002	0	*	*	*	*	*	*
2001	10,360	34.4	23.8	35.9	0.3	1.6	4.1
2000	0	*	*	*	*	*	*
1999	0	*	*	*	*	*	*
1998	0	*	*	*	*	*	*
1997	0	*	*	*	*	*	*
1996	0	*	*	*	*	*	*
1995	0	*	*	*	*	*	*
1994	0	*	*	*	*	*	*
1993	0	*	*	*	*	*	*
1992	0	*	*	*	*	*	*
1991	0	*	*	*	*	*	*
1990	0	*	*	*	*	*	*
1989	409	48.4	32.0	19.1	0.5	0.0	0.0
1988	812	46.7	30.4	22.0	0.5	0.4	0.0
1987	555	49.2	31.5	18.9	0.2	0.2	0.0
1986	331	58.0	29.0	12.7	0.3	0.0	0.0
1985	545	51.0	31.0	16.5	1.3	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 14C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Racial and Ethnic Distribution**

<b>Summary of Trends in Racial and Ethnic Distribution</b> <sup>(2)</sup>							
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
1984	534	58.1	29.0	11.8	0.9	0.2	0.0
1983	430	58.8	26.7	13.7	0.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 15C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Age Distribution**

<b>Summary of Trends in Age Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2005	10,624	0.3	6.5	11.4	57.5	22.3	1.9	0.0
2004	10,219	0.4	7.1	12.0	57.0	21.6	1.8	0.0
2003	14,142	0.5	6.7	12.5	57.4	21.3	1.6	0.0
2002	0	*	*	*	*	*	*	*
2001	10,360	0.5	7.2	12.1	54.9	23.4	1.9	0.0
2000	0	*	*	*	*	*	*	*
1999	0	*	*	*	*	*	*	*
1998	0	*	*	*	*	*	*	*
1997	0	*	*	*	*	*	*	*
1996	0	*	*	*	*	*	*	*
1995	0	*	*	*	*	*	*	*
1994	0	*	*	*	*	*	*	*
1993	0	*	*	*	*	*	*	*
1992	0	*	*	*	*	*	*	*
1991	0	*	*	*	*	*	*	*
1990	0	*	*	*	*	*	*	*
1989	409	0.5	10.0	14.4	57.7	17.1	0.2	0.0
1988	812	0.4	12.3	14.0	56.7	15.5	1.1	0.0
1987	555	0.7	8.1	12.6	60.0	17.8	0.7	0.0
1986	331	0.9	12.1	16.3	58.3	12.1	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 15C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Age Distribution**

<b>Summary of Trends in Age Distribution</b> <sup>(2)</sup>								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1985	545	1.5	12.8	17.6	51.6	15.6	0.9	0.0
1984	534	0.6	12.0	17.6	56.4	12.4	1.1	0.0
1983	430	0.5	12.8	16.3	59.3	10.9	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 16C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Health Indicators**

<b>Summary of Trends in Maternal Health Indicators</b> <sup>(2)</sup>										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2005	10,536	9.5	44.2	10,066	69.7	12.9	1,956	23.2	7,271	58.1
2004	10,137	11.2	42.6	9,701	68.9	14.1	1,939	26.7	7,130	59.2
2003	13,788	11.7	40.9	10,795	24.8	46.0	2,919	26.6	8,059	57.6
2002	0	*	*	0	*	*	0	*	0	*
2001	10,215	11.8	41.0	10,171	26.7	43.3	1,987	29.9	7,737	57.9
2000	0	*	*	0	*	*	0	*	0	*
1999	0	*	*	0	*	*	0	*	0	*
1998	0	*	*	0	*	*	0	*	0	*
1997	0	*	*	0	*	*	0	*	0	*
1996	0	*	*	0	*	*	0	*	0	*
1995	0	*	*	0	*	*	0	*	0	*
1994	0	*	*	0	*	*	0	*	0	*
1993	0	*	*	0	*	*	0	*	0	*
1992	0	*	*	0	*	*	0	*	0	*
1991	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 16C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Health Indicators**

<b>Summary of Trends in Maternal Health Indicators</b> <sup>(2)</sup>										
Year	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1990	0	*	*	0	*	*	0	*	0	*
1989	408	20.1	23.5	0	*	*	99	*	136	34.6
1988	789	18.0	28.3	0	*	*	132	22.7	152	39.5
1987	549	19.7	26.8	0	*	*	134	34.3	126	24.6
1986	325	19.4	24.9	0	*	*	67	*	107	23.4
1985	541	22.0	22.4	0	*	*	118	19.5	214	20.1
1984	533	22.7	18.4	0	*	*	126	25.4	255	21.2
1983	426	20.7	21.4	0	*	*	96	*	217	26.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 17C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Behavioral Indicators**

<b>Summary of Trends in Maternal Behavioral Indicators</b> <sup>(2)</sup>					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2005	10,135	86.7	2.8	10,264	21.9
2004	9,764	84.7	3.7	9,718	20.3
2003	11,377	84.9	4.4	13,888	17.8
2002	0	*	*	0	*
2001	10,331	81.8	6.8	10,160	17.9
2000	0	*	*	0	*
1999	0	*	*	0	*
1998	0	*	*	0	*
1997	0	*	*	0	*
1996	0	*	*	0	*
1995	0	*	*	0	*
1994	0	*	*	0	*
1993	0	*	*	0	*
1992	0	*	*	0	*
1991	0	*	*	0	*
1990	0	*	*	0	*
1989	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 17C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**  
**Summary of Trends in Maternal Behavioral Indicators**

<b>Summary of Trends in Maternal Behavioral Indicators</b> <sup>(2)</sup>					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
1988	0	*	*	0	*
1987	0	*	*	0	*
1986	0	*	*	0	*
1985	0	*	*	0	*
1984	0	*	*	0	*
1983	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 18C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators**

<b>Summary of Trends in Smoking Indicators</b> (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
2005	10,605	27.3	10,142	3.7	2,358	53.7	9,093	16.9	10,142	9.6
2004	2,810	99.8	9,773	4.5	2,301	51.1	8,706	18.5	9,771	10.9
2003	14,099	26.0	11,385	15.6	2,255	42.0	11,431	20.8	10,929	11.8
2002	0	*	0	*	0	*	0	*	0	*
2001	10,320	27.1	10,346	15.3	2,308	37.9	8,079	24.5	10,271	14.8
2000	0	*	0	*	0	*	0	*	0	*
1999	0	*	0	*	0	*	0	*	0	*
1998	0	*	0	*	0	*	0	*	0	*
1997	0	*	0	*	0	*	0	*	0	*
1996	0	*	0	*	0	*	0	*	0	*
1995	0	*	0	*	0	*	0	*	0	*
1994	0	*	0	*	0	*	0	*	0	*
1993	0	*	0	*	0	*	0	*	0	*
1992	0	*	0	*	0	*	0	*	0	*
1991	0	*	0	*	0	*	0	*	0	*
1990	0	*	0	*	0	*	0	*	0	*
1989	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 18C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators**

<b>Summary of Trends in Smoking Indicators</b> (2)									
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum	
	Number	%	Number	%		Number	%	Number	%
1988	0	*	0	*	0	*	0	*	
1987	0	*	0	*	0	*	0	*	
1986	0	*	0	*	0	*	0	*	
1985	0	*	0	*	0	*	0	*	
1984	0	*	0	*	0	*	0	*	
1983	0	*	0	*	0	*	0	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 19C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Infant Health Indicators**

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2005	10,126	7.7	7.6	9,763	8.9	8,888	3.5	10,137	61.9
2004	9,757	7.2	7.4	9,267	7.8	8,540	3.4	9,763	59.7
2003	9,747	7.4	7.6	10,668	9.0	8,767	3.7	10,791	42.1
2002	0	*	*	0	*	0	*	0	*
2001	9,016	7.1	8.6	9,985	8.8	8,113	3.5	10,093	40.0
2000	0	*	*	0	*	0	*	0	*
1999	0	*	*	0	*	0	*	0	*
1998	0	*	*	0	*	0	*	0	*
1997	0	*	*	0	*	0	*	0	*
1996	0	*	*	0	*	0	*	0	*
1995	0	*	*	0	*	0	*	0	*
1994	0	*	*	0	*	0	*	0	*
1993	0	*	*	0	*	0	*	0	*
1992	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 19C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Infant Health Indicators**

Summary of Trends in Infant Health Indicators (2)									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1991	0	*	*	0	*	0	*	0	*
1990	0	*	*	0	*	0	*	0	*
1989	409	7.6	7.1	400	0.5	398	7.5	100	100.0
1988	411	5.8	11.4	409	0.7	406	5.7	95	*
1987	354	6.2	9.6	350	0.9	347	6.1	63	*
1986	324	7.4	11.1	322	1.6	317	7.3	65	*
1985	525	6.1	9.3	522	0.8	518	6.0	99	*
1984	500	9.2	9.4	495	2.4	483	8.9	149	100.0
1983	424	7.3	6.4	422	4.0	405	6.7	123	100.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2005	3,466	11.1	44.1	3,351	62.8	18.8	644	17.9	2,102	54.7
	2004	3,351	13.3	42.1	3,254	61.6	19.2	645	21.1	2,106	57.2
	2003	4,361	13.4	41.2	3,585	20.5	52.5	908	21.5	2,336	54.0
	2002	0	*	*	0	*	*	0	*	0	*
	2001	3,527	14.1	41.1	3,510	23.0	47.2	673	24.8	2,373	55.0
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
Black, Not Hispanic	2005	2,314	7.4	51.4	2,181	70.8	12.1	497	31.8	1,711	62.2
	2004	2,340	9.1	51.9	2,206	68.3	14.1	522	36.2	1,786	64.6
	2003	3,377	9.4	47.9	2,490	26.5	45.7	855	34.4	2,041	63.4
	2002	0	*	*	0	*	*	0	*	0	*
	2001	2,430	8.7	50.7	2,413	28.9	43.3	565	38.2	2,019	63.6
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2005	4,442	8.8	41.8	4,228	74.3	8.9	757	21.8	3,254	58.5
	2004	3,928	9.5	39.3	3,733	75.1	10.2	639	24.9	2,858	58.0
	2003	5,315	10.9	38.0	4,091	26.5	42.1	994	25.7	3,210	57.1
	2002	0	*	*	0	*	*	0	*	0	*
	2001	3,673	10.5	37.2	3,663	27.7	40.9	638	29.3	2,900	57.4
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
American Indian/Alaskan Native	2005	45	*	*	41	*	*	8	*	27	*
	2004	22	*	*	20	*	*	6	*	19	*
	2003	21	*	*	15	*	*	5	*	10	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	26	*	*	26	*	*	7	*	21	*
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2005	223	18.8	25.6	219	79.5	7.3	43	*	147	53.1
	2004	178	30.3	15.7	176	78.4	6.8	48	*	125	48.8
	2003	214	31.8	15.9	186	37.1	26.3	56	*	127	52.8
	2002	0	*	*	0	*	*	0	*	0	*
	2001	166	28.9	15.1	166	30.7	34.3	31	*	119	55.5
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*
All Other/Unknown	2005	46	*	*	46	*	*	7	*	30	*
	2004	318	14.5	36.2	312	70.5	12.8	79	*	236	54.7
	2003	500	13.0	31.8	428	28.5	40.2	101	22.8	335	53.1
	2002	0	*	*	0	*	*	0	*	0	*
	2001	393	15.3	27.2	393	34.4	36.6	73	*	305	49.5
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 20C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**

**Summary of Trends in Maternal Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Health Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Total All Race/Ethnic Groups	2005	10,536	9.5	44.2	10,066	69.7	12.9	1,956	23.2	7,271	58.1
	2004	10,137	11.2	42.6	9,701	68.9	14.1	1,939	26.7	7,130	59.2
	2003	13,788	11.7	40.9	10,795	24.8	46.0	2,919	26.6	8,059	57.6
	2002	0	*	*	0	*	*	0	*	0	*
	2001	10,215	11.8	41.0	10,171	26.7	43.3	1,987	29.9	7,737	57.9
	2000	0	*	*	0	*	*	0	*	0	*
	1999	0	*	*	0	*	*	0	*	0	*
	1998	0	*	*	0	*	*	0	*	0	*
	1997	0	*	*	0	*	*	0	*	0	*
	1996	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2005	3,385	88.5	3.4	3,378	23.0
	2004	3,281	87.1	4.0	3,231	20.4
	2003	3,732	87.6	4.6	4,373	17.9
	2002	0	*	*	0	*
	2001	3,551	86.3	5.7	3,488	18.1
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
Black, Not Hispanic	2005	2,190	84.4	2.5	2,270	19.4
	2004	2,219	83.0	3.4	2,255	20.8
	2003	2,604	82.6	4.2	3,376	15.8
	2002	0	*	*	0	*
	2001	2,456	79.2	7.0	2,415	16.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2005	4,250	86.8	2.3	4,315	22.7
	2004	3,743	84.7	3.0	3,717	21.1
	2003	4,336	84.7	3.7	5,359	19.7
	2002	0	*	*	0	*
	2001	3,708	80.3	7.0	3,652	19.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
American Indian/Alaskan Native	2005	41	*	*	42	*
	2004	20	*	*	21	*
	2003	16	*	*	21	*
	2002	0	*	*	0	*
	2001	26	*	*	26	*
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2005	223	78.5	6.3	213	14.1
	2004	181	78.5	6.1	171	14.0
	2003	192	80.2	5.7	214	8.9
	2002	0	*	*	0	*
	2001	168	78.0	10.7	165	11.5
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*
All Other/Unknown	2005	46	*	*	46	*
	2004	320	75.9	8.4	323	11.1
	2003	497	80.1	8.7	545	13.9
	2002	0	*	*	0	*
	2001	422	74.4	12.3	414	11.6
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 21C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity**

<b>Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity</b> <sup>(2)</sup>						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Total All Race/Ethnic Groups	2005	10,135	86.7	2.8	10,264	21.9
	2004	9,764	84.7	3.7	9,718	20.3
	2003	11,377	84.9	4.4	13,888	17.8
	2002	0	*	*	0	*
	2001	10,331	81.8	6.8	10,160	17.9
	2000	0	*	*	0	*
	1999	0	*	*	0	*
	1998	0	*	*	0	*
	1997	0	*	*	0	*
	1996	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2005	3,508	45.3	3,390	8.3	1,240	47.3	2,739	26.5	3,390	15.8
	2004	1,601	100.0	3,285	9.9	1,250	44.6	2,625	30.5	3,285	17.4
	2003	4,447	45.2	3,734	25.9	1,243	37.7	3,352	32.7	3,637	19.1
	2002	0	*	0	*	0	*	0	*	0	*
	2001	3,550	46.8	3,556	28.3	1,306	34.3	2,569	38.3	3,540	23.2
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
Black, Not Hispanic	2005	2,319	23.6	2,191	2.0	475	54.9	2,110	16.5	2,191	7.9
	2004	524	99.6	2,223	2.4	449	53.9	2,121	16.1	2,222	8.0
	2003	3,406	20.8	2,606	13.0	430	40.9	2,894	19.5	2,517	9.2
	2002	0	*	0	*	0	*	0	*	0	*
	2001	2,447	21.8	2,462	12.1	481	37.4	2,005	20.5	2,450	11.6
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2005	4,460	16.1	4,251	1.1	614	65.3	4,002	10.6	4,251	5.6
	2004	618	99.4	3,744	1.4	553	62.0	3,573	11.6	3,743	7.1
	2003	5,441	16.1	4,339	9.6	547	50.8	4,680	13.7	4,127	7.6
	2002	0	*	0	*	0	*	0	*	0	*
	2001	3,708	14.7	3,712	6.8	486	47.3	3,108	16.7	3,688	9.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
American Indian/Alaskan Native	2005	45	*	41	*	8	*	34	*	41	*
	2004	5	*	20	*	3	*	19	*	20	*
	2003	22	*	16	*	0	*	15	*	15	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	26	*	26	*	2	*	20	*	26	*
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2005	227	5.3	223	0.4	11	*	173	12.1	223	9.9
	2004	17	*	181	0.0	13	*	140	12.1	181	6.6
	2003	219	7.3	193	2.6	12	*	150	18.0	190	4.7
	2002	0	*	0	*	0	*	0	*	0	*
	2001	168	6.5	168	2.4	9	*	109	17.4	168	12.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*
All Other/Unknown	2005	46	*	46	*	10	*	35	*	46	*
	2004	45	*	320	2.2	33	*	228	15.8	320	10.3
	2003	564	10.5	497	8.9	23	*	340	14.4	443	8.1
	2002	0	*	0	*	0	*	0	*	0	*
	2001	421	10.0	422	6.4	24	*	268	15.7	399	8.5
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 22C**  
**2005 Pregnancy Nutrition Surveillance** (1)  
**Connecticut**  
**Summary of Trends in Smoking Indicators by Race/Ethnicity**

<b>Summary of Trends in Smoking Indicators by Race/Ethnicity</b> (2)											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Total All Race/Ethnic Groups	2005	10,605	27.3	10,142	3.7	2,358	53.7	9,093	16.9	10,142	9.6
	2004	2,810	99.8	9,773	4.5	2,301	51.1	8,706	18.5	9,771	10.9
	2003	14,099	26.0	11,385	15.6	2,255	42.0	11,431	20.8	10,929	11.8
	2002	0	*	0	*	0	*	0	*	0	*
	2001	10,320	27.1	10,346	15.3	2,308	37.9	8,079	24.5	10,271	14.8
	2000	0	*	0	*	0	*	0	*	0	*
	1999	0	*	0	*	0	*	0	*	0	*
	1998	0	*	0	*	0	*	0	*	0	*
	1997	0	*	0	*	0	*	0	*	0	*
	1996	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2005	3,385	7.1	9.1	3,235	9.1	2,940	3.1	3,387	56.3
	2004	3,280	6.2	8.8	3,118	8.0	2,868	2.6	3,282	53.3
	2003	3,236	6.2	9.5	3,507	9.1	2,899	2.9	3,574	32.8
	2002	0	*	*	0	*	0	*	0	*
	2001	3,150	6.0	10.2	3,419	8.2	2,845	2.8	3,470	31.5
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
Black, Not Hispanic	2005	2,185	10.9	6.1	2,137	10.5	1,909	4.8	2,190	54.4
	2004	2,218	9.7	5.5	2,114	8.8	1,926	5.3	2,221	53.2
	2003	2,177	9.9	6.0	2,435	10.3	1,931	4.8	2,465	40.2
	2002	0	*	*	0	*	0	*	0	*
	2001	2,103	9.9	6.3	2,365	9.7	1,867	5.2	2,387	35.7
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2005	4,246	6.5	7.5	4,100	8.0	3,770	3.0	4,250	69.5
	2004	3,739	6.4	7.1	3,531	7.0	3,283	3.0	3,740	67.6
	2003	3,776	7.0	7.2	4,080	8.6	3,426	3.8	4,097	48.9
	2002	0	*	*	0	*	0	*	0	*
	2001	3,251	6.2	8.9	3,603	8.9	2,941	3.1	3,634	48.0
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
American Indian/Alaskan Native	2005	41	*	*	36	*	33	*	41	*
	2004	20	*	*	19	*	17	*	20	*
	2003	11	*	*	12	*	9	*	12	*
	2002	0	*	*	0	*	0	*	0	*
	2001	22	*	*	26	*	18	*	25	*
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)	High (4)							
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2005	223	9.0	4.5	209	7.7	193	5.2	223	73.5
	2004	180	10.0	6.1	167	9.0	152	2.6	180	67.8
	2003	170	9.4	0.6	183	7.7	155	3.9	188	47.9
	2002	0	*	*	0	*	0	*	0	*
	2001	140	7.1	5.7	163	9.2	125	2.4	164	42.7
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*
All Other/Unknown	2005	46	*	*	46	*	43	*	46	*
	2004	320	7.5	9.7	318	7.5	294	3.1	320	72.8
	2003	377	5.3	8.2	451	7.1	347	2.6	455	62.0
	2002	0	*	*	0	*	0	*	0	*
	2001	350	7.1	7.4	409	8.1	317	3.2	413	62.2
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 23C**  
**2005 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Connecticut**

**Summary of Trends in Infant Health Indicators by Race/Ethnicity**

<b>Summary of Trends in Infant Health Indicators by Race/Ethnicity</b> <sup>(2)</sup>										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Total All Race/Ethnic Groups	2005	10,126	7.7	7.6	9,763	8.9	8,888	3.5	10,137	61.9
	2004	9,757	7.2	7.4	9,267	7.8	8,540	3.4	9,763	59.7
	2003	9,747	7.4	7.6	10,668	9.0	8,767	3.7	10,791	42.1
	2002	0	*	*	0	*	0	*	0	*
	2001	9,016	7.1	8.6	9,985	8.8	8,113	3.5	10,093	40.0
	2000	0	*	*	0	*	0	*	0	*
	1999	0	*	*	0	*	0	*	0	*
	1998	0	*	*	0	*	0	*	0	*
	1997	0	*	*	0	*	0	*	0	*
	1996	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.