

Table 1D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Demographic Indicators
Children Aged < 5 Years (2,3)

Total Records	Number
Individual Children	7,599,318
Total Records (All Visits)	11,070,513

Source of Data	Number	%
WIC	6,322,267	83.5
EPSDT	464,456	6.1
Child Health-MCH	43,032	0.6
Child Health-HMO	740,032	9.8
Headstart	0	0.0
School Health	0	0.0
Other	0	0.0
Total	7,569,787	100.0
<i>Excluded Unknown:</i> 29,531 0.4 %		

Race/Ethnic Distribution	Number	%
White, Not Hispanic	2,397,477	32.2
Black, Not Hispanic	1,403,725	18.9
Hispanic	3,121,887	42.0
American Indian/Alaskan Native	77,766	1.0
Asian/Pacific Islander	190,710	2.6
Multiple Races	114,485	1.5
All Other	130,633	1.8
Total	7,436,683	100.0
<i>Excluded Unknown:</i> 162,635 2.1 %		

Age Distribution (4)	Number	%
0 - 5 Months	2,161,176	28.4
6 - 11 Months	696,262	9.2
12 - 23 Months	1,655,488	21.8
24 - 35 Months	1,156,306	15.2
3 - 4 Years	1,930,086	25.4
Total	7,599,318	100.0

% of Poverty Level	Number	%
0-50	1,391,840	31.0
51-100	1,260,627	28.1
101-130	541,072	12.0
131-150	255,595	5.7
151-185	313,858	7.0
186-200	17,508	0.4
Over 200	40,995	0.9
Adjunctive Eligibility	669,567	14.9
Total	4,491,062	100.0
<i>Excluded Unknown:</i> 3,108,256 40.9 %		

Program Participation	Number	%
WIC	4,540,760	98.8
FoodStamps	4,284,241	21.0
Medicaid	4,507,788	65.6
TANF	4,264,114	5.6
<i>Excluded Unknown:</i>		
WIC	3,058,558	40.2
FoodStamps	3,315,077	43.6
Medicaid	3,091,530	40.7
TANF	3,335,204	43.9

Migrant Status	Number	%
Yes	19,613	0.5
No	4,307,655	99.5
Total	4,327,268	100.0
<i>Excluded Unknown:</i> 3,272,050 43.1 %		

(1) Reporting period is January 1 through December 31.
 (2) Analyses based on one record per child.
 (3) Excludes records with unknown data or errors.
 (4) Records rejected if date of birth and date of visit unknown.
 * Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 2D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Summary of Health Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Health Indicators ⁽³⁾			
		National Prevalence	
		Number	%
Birthweight ⁽⁴⁾			
Low	< 2500 g	1,842,970	9.2
High	> 4000 g	1,842,970	6.6
Height and Weight			
Short Stature ⁽⁵⁾			
	< 5th	7,192,043	6.4
	< 10th	7,192,043	11.4
Underweight ⁽⁶⁾			
	< 5th	7,192,043	4.7
	< 10th	7,192,043	8.0
Overweight ⁽⁶⁾			
	≥ 95th	7,192,043	13.8
≥ 2 Yrs Overweight ⁽⁶⁾			
	85th - < 95th	2,945,212	16.4
	≥ 95th	2,945,212	14.8
Anemia ⁽⁷⁾			
Low Hb		4,092,770	14.3
Low Hct		652,304	10.9
Low Hb/Hct		4,521,591	14.0
Breastfeeding ⁽⁸⁾			
Ever Breastfed		1,577,832	60.1
Breastfed At Least 6 Months		790,341	25.2
Breastfed At Least 12 Months		875,205	18.1
Smoking in Household			
≤ 2 Hours TV Viewing/ Day ⁽⁹⁾		2,855,916	10.6
		584,732	73.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(9) Children aged 2 years and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 3D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Summary of Breastfeeding Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Breastfeeding Indicators ⁽³⁾		
	National Prevalence	
	Number	%
Breastfeeding Duration ^(4,5)		
Ever Breastfed	1,577,832	60.1
Breastfed 1+ Weeks	1,299,520	58.5
Breastfed 2+ Weeks	1,030,249	55.2
Breastfed 3+ Weeks	852,030	52.5
Breastfed 4+ Weeks	746,660	49.9
Breastfed 6+ Weeks	628,208	44.0
Breastfed 2+ Months	581,374	38.7
Breastfed 3+ Months	610,380	33.8
Breastfed 4+ Months	683,087	29.9
Breastfed 5+ Months	763,521	27.0
Breastfed 6+ Months	790,341	25.2
Breastfed 9+ Months	820,087	20.7
Breastfed 12+ Months	875,205	18.1
Breastfed 18+ Months	660,721	12.2
Exclusive Breastfeeding ⁽⁶⁾		
Exclusively Breastfed at Least 3 Months	104,492	10.3
Exclusively Breastfed at Least 6 Months	143,261	5.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. For each breastfeeding duration category, the analysis includes only children who turned that age during the reporting period by/on their date of visit. For example, infants who turned 9 months old during the reporting period by/on their date of visit are included in the 9+ months analysis.

(5) It is possible the analysis will show that slightly more infants are breastfed at older ages, e.g. 1+ week of age, than at younger ages, e.g. at birth (ever breastfed), due to variation in infants included in each breastfeeding analysis category.

(6) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Comparison of Racial and Ethnic Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Contributor								
Contributor	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Alabama	130,882	43.6	43.1	10.6	0.4	0.1	1.7	0.6
Arizona	153,725	21.1	4.5	70.5	1.7	1.0	1.2	0.0
Arkansas	74,669	55.6	26.0	15.8	0.6	1.1	0.9	0.0
AZ I/Tribal Council	12,121	2.1	0.4	30.1	65.4	0.2	1.7	0.0
California	1,240,441	8.2	4.8	67.6	0.3	4.0	0.0	15.2
Cheyenne River-SD	762	0.5	0.0	2.9	53.3	0.0	43.3	0.0
Colorado	97,607	29.1	5.9	59.4	1.0	1.4	3.2	0.1
Connecticut	55,013	26.7	24.8	45.3	0.4	1.8	0.9	0.0
D.C.	13,667	1.9	70.4	24.8	0.2	2.4	0.2	0.0
Florida	434,873	27.1	29.7	40.1	0.2	0.9	2.1	0.0
Georgia	281,431	34.1	42.0	19.2	0.5	4.1	0.0	0.0
Idaho	40,736	63.3	1.0	31.4	2.1	1.0	1.2	0.0
Illinois	275,811	29.6	24.7	40.9	0.6	2.0	0.0	2.1
Indiana	144,820	58.2	17.7	19.0	0.2	0.9	4.0	0.0
Iowa	72,143	61.6	7.7	22.8	0.1	1.3	3.8	2.7
Kansas	53,321	52.7	10.1	31.2	0.9	1.6	3.5	0.0
Kentucky	135,754	78.7	11.7	7.2	0.1	0.9	1.3	0.0
Maryland	112,865	22.8	47.2	23.0	0.3	2.6	4.2	0.0
Massachusetts	122,006	42.8	19.2	32.3	0.1	5.6	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Comparison of Racial and Ethnic Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Contributor								
Contributor	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Michigan	232,137	53.1	26.5	13.7	0.4	1.6	4.7	0.0
Minnesota	129,890	47.5	15.9	19.3	3.4	7.4	6.2	0.3
Missouri	138,503	66.0	20.8	10.8	0.3	1.1	0.0	1.0
Montana	20,701	64.7	0.4	7.1	15.2	0.3	12.3	0.0
Navajo Nation	14,162	0.7	0.1	1.8	97.4	0.1	0.0	0.0
Nebraska	45,148	48.8	11.0	34.2	1.5	1.4	3.1	0.0
Nevada	51,271	18.0	8.5	66.0	0.4	2.4	4.7	0.0
New Hampshire	17,135	87.4	4.6	5.0	0.4	1.6	0.9	0.0
New Jersey	157,001	17.4	26.3	51.3	0.5	3.3	1.0	0.1
New Mexico	66,228	15.6	1.6	76.6	3.5	0.5	1.1	1.1
New York	467,520	28.3	24.7	37.5	0.7	7.3	1.4	0.1
North Carolina	211,961	37.2	31.9	28.0	1.1	1.2	0.0	0.5
Ohio Child Hlth	27,757	36.8	40.5	14.9	0.1	1.2	2.8	3.7
Ohio-WIC	281,596	59.6	30.6	7.9	0.5	1.3	0.0	0.0
Oregon	100,604	50.2	2.8	40.1	1.6	2.6	2.2	0.5
Pennsylvania	244,213	53.1	22.9	19.2	0.3	2.5	2.0	0.0
Puerto Rico	198,967	0.0	0.0	58.2	0.0	0.0	0.0	41.8
Rhode Island	23,875	42.3	14.4	39.0	0.5	2.2	1.6	0.0
Rosebud Sioux-SD	1,359	1.6	0.0	2.1	89.8	0.0	6.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Racial and Ethnic Distribution by Contributor
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution by Contributor								
Contributor	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
South Carolina	100,602	37.0	47.7	12.5	0.1	0.2	0.0	2.5
South Dakota	20,384	60.1	2.8	6.6	26.7	0.8	2.9	0.1
Standing Rock-ND	982	0.9	0.0	0.4	97.5	0.0	1.2	0.0
Tennessee	156,123	58.6	26.5	13.0	0.0	0.0	0.0	1.8
Texas	922,732	12.6	11.2	73.9	0.1	1.0	1.3	0.0
Vermont	13,265	93.7	2.6	0.2	0.5	1.0	0.0	1.9
Virginia	135,114	38.7	37.8	18.9	0.3	2.5	1.7	0.0
Washington	193,741	42.3	6.4	35.6	3.9	4.9	6.4	0.4
West Virginia	52,027	88.0	3.7	2.3	0.0	0.2	5.7	0.0
Wisconsin	121,673	45.1	18.6	24.2	2.0	4.5	5.1	0.5
Nation	7,599,318	31.5	18.5	41.1	1.0	2.5	1.5	3.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Age Distribution by Contributor
Children Aged < 5 Years (2)

Comparison of Age Distribution by Contributor				
Contributor	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Alabama	130,882	37.3	23.7	39.0
Arizona	153,725	39.2	23.4	37.4
Arkansas	74,669	38.0	22.6	39.4
AZ I/Tribal Council	12,121	29.4	25.6	45.0
California	1,240,441	46.2	20.6	33.2
Cheyenne River-SD	762	28.6	22.0	49.3
Colorado	97,607	42.6	18.9	38.5
Connecticut	55,013	31.6	25.3	43.1
D.C.	13,667	38.5	23.5	38.1
Florida	434,873	39.5	21.4	39.1
Georgia	281,431	44.1	19.4	36.5
Idaho	40,736	37.4	20.4	42.3
Illinois	275,811	43.8	19.6	36.5
Indiana	144,820	40.7	21.0	38.2
Iowa	72,143	38.9	18.4	42.7
Kansas	53,321	37.2	20.3	42.5
Kentucky	135,754	40.5	18.7	40.8
Maryland	112,865	37.7	23.1	39.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Age Distribution by Contributor
Children Aged < 5 Years (2)

Comparison of Age Distribution by Contributor				
Contributor	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Massachusetts	122,006	33.5	20.5	46.0
Michigan	232,137	36.7	21.2	42.1
Minnesota	129,890	28.9	25.7	45.4
Missouri	138,503	34.3	26.1	39.6
Montana	20,701	34.3	22.2	43.5
Navajo Nation	14,162	26.6	24.7	48.7
Nebraska	45,148	38.0	19.7	42.2
Nevada	51,271	44.1	19.0	36.8
New Hampshire	17,135	36.5	21.5	42.0
New Jersey	157,001	40.4	19.6	40.0
New Mexico	66,228	35.9	20.8	43.3
New York	467,520	30.5	25.2	44.3
North Carolina	211,961	36.1	21.7	42.2
Ohio Child Hlth	27,757	37.6	21.1	41.4
Ohio-WIC	281,596	41.2	17.8	41.0
Oregon	100,604	35.0	19.1	46.0
Pennsylvania	244,213	33.0	23.8	43.2
Puerto Rico	198,967	28.2	21.6	50.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Age Distribution by Contributor
Children Aged < 5 Years (2)

Comparison of Age Distribution by Contributor				
Contributor	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Rhode Island	23,875	33.2	20.9	45.9
Rosebud Sioux-SD	1,359	28.2	20.2	51.7
South Carolina	100,602	42.4	22.9	34.7
South Dakota	20,384	40.9	17.8	41.4
Standing Rock-ND	982	24.1	23.0	52.9
Tennessee	156,123	35.2	24.3	40.5
Texas	922,732	29.5	24.4	46.1
Vermont	13,265	28.5	19.7	51.8
Virginia	135,114	30.3	26.8	42.8
Washington	193,741	36.7	18.7	44.6
West Virginia	52,027	42.0	16.8	41.3
Wisconsin	121,673	36.7	19.9	43.3
Nation	7,599,318	37.6	21.8	40.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alabama	37,278	12.4 (45)	4.9 (3)	125,927	6.8 (31)	7.5 (46)	11.8 (9)	49,826	15.0 (8)	13.7 (17)	80,207	19.0 (44)
Arizona	41,569	8.4 (15)	6.9 (29)	150,446	7.8 (41)	3.4 (17)	14.8 (32)	56,405	15.1 (9)	13.5 (15)	92,485	16.5 (38)
Arkansas	25,163	9.7 (29)	5.9 (10)	71,325	8.2 (43)	4.9 (32)	10.8 (5)	28,481	15.2 (10)	13.2 (13)	40,922	15.7 (32)
AZ I/Tribal Council	2,672	7.7 (8)	9.2 (43)	10,729	7.0 (33)	2.0 (3)	23.3 (48)	5,203	20.8 (45)	24.1 (47)	6,886	10.4 (12)
California	229,856	6.4 (2)	7.7 (36)	1,056,397	5.1 (9)	5.5 (36)	15.4 (39)	339,961	16.2 (19)	17.0 (39)	527,868	14.0 (29)
Cheyenne River-SD	156	8.3 (11)	7.7 (36)	723	1.5 (1)	1.4 (2)	23.0 (46)	353	23.2 (46)	18.4 (44)	454	16.1 (35)
Colorado	26,634	10.1 (36)	4.6 (1)	96,600	8.1 (42)	3.7 (21)	8.2 (1)	37,111	14.7 (5)	9.6 (1)	51,974	12.1 (19)
Connecticut	15,230	9.5 (27)	6.7 (24)	53,416	3.9 (4)	7.7 (47)	12.2 (14)	22,854	16.5 (25)	16.2 (35)	36,481	9.4 (8)
D.C.	4,508	12.0 (44)	4.9 (3)	13,067	9.1 (45)	4.7 (31)	13.1 (24)	5,107	14.8 (7)	15.4 (31)	7,697	28.8 (47)
Florida	123,782	9.7 (29)	5.9 (10)	424,890	4.8 (6)	5.3 (33)	12.7 (21)	166,498	15.4 (11)	13.9 (20)	259,805	16.7 (40)
Georgia	80,245	10.4 (38)	5.4 (6)	275,377	7.3 (38)	3.6 (20)	13.6 (28)	101,052	16.4 (23)	14.5 (27)	163,053	13.2 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Idaho	10,530	7.5 (7)	7.5 (31)	38,127	7.0 (33)	4.0 (24)	10.2 (2)	16,305	16.9 (28)	12.4 (6)	26,006	12.7 (23)
Illinois	84,621	9.7 (29)	6.2 (16)	269,173	7.4 (39)	5.3 (33)	12.6 (19)	98,910	15.4 (11)	14.4 (26)	147,984	10.9 (15)
Indiana	42,199	9.5 (27)	6.3 (18)	141,229	6.7 (28)	3.7 (21)	11.6 (8)	54,591	16.4 (23)	14.0 (21)	93,853	15.9 (33)
Iowa	17,935	8.8 (20)	7.7 (36)	70,491	5.8 (19)	2.8 (7)	13.9 (29)	30,161	17.9 (37)	14.6 (28)	43,572	9.4 (8)
Kansas	8,310	8.3 (11)	6.7 (24)	52,465	6.2 (23)	2.5 (5)	13.4 (26)	22,404	17.0 (31)	13.8 (19)	35,746	9.4 (8)
Kentucky	35,259	10.2 (37)	6.1 (14)	129,341	7.0 (33)	5.3 (33)	15.3 (38)	53,777	17.9 (37)	17.4 (41)	84,401	12.3 (21)
Maryland	28,622	11.2 (43)	6.0 (12)	106,452	6.6 (24)	3.4 (17)	13.3 (25)	43,428	16.3 (21)	14.8 (29)	66,435	22.3 (46)
Massachusetts	30,052	9.0 (23)	7.5 (31)	118,367	4.8 (6)	5.6 (37)	14.8 (32)	54,617	17.3 (33)	16.7 (38)	65,496	12.4 (22)
Michigan	55,559	9.8 (33)	7.3 (30)	225,665	6.8 (31)	4.1 (26)	12.4 (15)	95,780	16.2 (19)	13.3 (14)	135,168	12.9 (24)
Minnesota	32,560	7.4 (5)	9.1 (42)	126,384	5.1 (9)	3.0 (10)	12.1 (12)	57,609	16.5 (25)	13.1 (10)	74,164	11.6 (18)
Missouri	42,993	9.1 (25)	6.6 (22)	134,102	7.1 (36)	4.5 (29)	11.8 (9)	53,585	16.6 (27)	13.6 (16)	85,027	17.5 (41)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Montana	5,004	8.1 (10)	7.6 (33)	20,457	5.2 (11)	2.9 (8)	12.7 (21)	9,031	16.1 (18)	12.1 (5)	15,710	9.1 (7)
Navajo Nation	3,046	7.2 (4)	6.5 (20)	13,824	5.5 (15)	1.1 (1)	15.1 (37)	6,787	20.5 (43)	16.3 (36)	10,758	8.1 (6)
Nebraska	11,507	8.6 (18)	6.7 (24)	44,248	5.7 (16)	4.5 (29)	12.6 (19)	18,708	15.7 (14)	13.1 (10)	27,519	16.4 (36)
Nevada	13,740	8.3 (11)	6.5 (20)	49,016	8.3 (44)	3.3 (15)	14.0 (30)	18,201	15.7 (14)	14.0 (21)	27,445	10.7 (14)
New Hampshire	4,039	8.0 (9)	9.0 (41)	15,460	6.7 (28)	3.1 (11)	15.0 (35)	7,003	18.1 (40)	15.9 (34)	9,085	13.8 (28)
New Jersey	42,986	9.3 (26)	6.2 (16)	153,295	6.1 (22)	4.1 (26)	16.3 (40)	61,327	17.5 (34)	18.1 (43)	101,537	18.9 (43)
New Mexico	15,928	10.0 (35)	4.6 (1)	64,109	7.7 (40)	6.1 (40)	10.5 (3)	28,089	14.0 (3)	11.5 (3)	39,895	10.0 (11)
New York	127,134	8.9 (22)	6.7 (24)	454,208	4.1 (5)	6.0 (39)	12.9 (23)	199,330	16.9 (28)	15.3 (30)	326,455	11.0 (16)
North Carolina	58,519	9.7 (29)	6.7 (24)	172,985	5.7 (16)	8.4 (48)	14.8 (32)	80,161	15.7 (14)	15.4 (31)	128,357	12.2 (20)
Ohio Child Hlth	0	* *	* *	25,854	9.6 (46)	4.4 (28)	16.8 (41)	10,642	17.1 (32)	17.4 (41)	0	* *
Ohio-WIC	68,490	11.1 (42)	5.7 (9)	276,483	6.6 (24)	4.0 (24)	12.1 (12)	114,008	15.5 (13)	11.7 (4)	188,701	13.6 (26)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2006 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Oregon	21,572	6.6 (3)	9.8 (45)	97,610	4.9 (8)	3.1 (11)	12.4 (15)	44,826	17.5 (34)	14.3 (24)	54,967	14.9 (31)
Pennsylvania	66,379	10.6 (39)	6.1 (14)	239,212	6.0 (21)	6.3 (42)	10.7 (4)	103,524	13.6 (1)	11.1 (2)	168,122	16.6 (39)
Puerto Rico	0	* *	* *	194,271	12.0 (47)	7.0 (45)	19.6 (44)	97,871	20.7 (44)	19.2 (45)	147,673	5.9 (2)
Rhode Island	4,511	8.8 (20)	7.6 (33)	21,812	6.6 (24)	3.4 (17)	17.8 (42)	10,449	17.9 (37)	16.5 (37)	14,871	16.4 (36)
Rosebud Sioux-SD	290	8.3 (11)	6.6 (22)	1,322	1.8 (3)	3.3 (15)	18.8 (43)	691	23.9 (47)	21.7 (46)	950	19.1 (45)
South Carolina	23,414	12.4 (45)	5.1 (5)	88,466	13.6 (48)	6.7 (44)	12.5 (18)	32,635	14.4 (4)	13.7 (17)	62,250	16.0 (34)
South Dakota	5,176	8.4 (15)	7.9 (39)	19,849	6.6 (24)	3.1 (11)	13.5 (27)	8,260	17.7 (36)	14.3 (24)	14,080	7.9 (5)
Standing Rock-ND	154	3.2 (1)	8.4 (40)	932	1.5 (1)	6.2 (41)	23.0 (46)	488	24.2 (48)	24.8 (48)	739	13.7 (27)
Tennessee	46,756	10.9 (41)	5.4 (6)	151,584	5.9 (20)	6.5 (43)	11.4 (7)	61,881	14.7 (5)	13.1 (10)	97,730	7.5 (4)
Texas	227,376	9.0 (23)	6.0 (12)	892,311	7.2 (37)	2.9 (8)	15.0 (35)	419,644	16.9 (28)	15.6 (33)	645,954	18.2 (42)
Vermont	2,357	8.5 (17)	10.8 (46)	12,847	5.3 (12)	3.1 (11)	12.0 (11)	6,697	15.8 (17)	12.9 (8)	9,442	6.1 (3)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2006 Pediatric Nutrition Surveillance (1)
National
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
Virginia	32,197	9.9 (34)	6.3 (18)	132,012	6.7 (28)	2.3 (4)	22.0 (45)	54,525	18.1 (40)	17.0 (39)	85,866	14.8 (30)
Washington	42,536	7.4 (5)	9.6 (44)	189,457	5.3 (12)	2.7 (6)	14.3 (31)	83,705	18.3 (42)	14.2 (23)	95,765	11.0 (16)
West Virginia	13,087	10.8 (40)	5.5 (8)	51,178	5.4 (14)	5.6 (37)	10.8 (5)	21,044	13.9 (2)	12.7 (7)	36,991	5.6 (1)
Wisconsin	31,039	8.6 (18)	7.6 (33)	118,548	5.7 (16)	3.8 (23)	12.4 (15)	51,667	16.3 (21)	13.0 (9)	85,045	10.4 (12)
Nation	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591	14.0

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7D
2006 Pediatric Nutrition Surveillance National (1)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4,5)

Contributor	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Alabama	39,433	33.9 (39)	13,922	2.0 (38)	0	* *	0	* *	0	* *	0	* *		
Arizona	41,495	61.1 (19)	31,581	25.1 (16)	27,478	11.7 (28)	12,459	8.1 (11)	15,004	2.7 (11)	0	* *	146,731 15.0 (10)	
Arkansas	24,892	45.2 (34)	2,597	9.2 (37)	8,551	5.2 (37)	0	* *	0	* *	0	* *	0	* *
AZ I/Tribal Council	2,946	60.2 (21)	1,050	31.7 (9)	2,021	27.1 (3)	549	12.4 (8)	949	7.9 (7)	5,112	69.1 (11)	10,532 9.6 (6)	
California	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *	1,235,658 4.6 (2)	
Cheyenne River-SD	85	* *	66	* *	62	* *	33	* *	52	* *	391	76.7 (5)	639 7.7 (4)	
Colorado	26,660	74.2 (6)	20,820	28.9 (12)	14,266	18.5 (12)	0	* *	0	* *	0	* *	0	* *
Connecticut	15,299	57.8 (25)	5,370	24.6 (17)	9,809	18.3 (13)	0	* *	0	* *	0	* *	0	* *
D.C.	0	* *	0	* *	0	* *	0	* *	0	* *	5,401	67.7 (12)	10,220 12.5 (8)	
Florida	123,859	66.3 (14)	80,866	24.4 (18)	72,223	14.4 (21)	0	* *	0	* *	0	* *	0	* *
Georgia	80,464	52.4 (31)	46,146	18.4 (29)	35,488	8.5 (33)	0	* *	0	* *	0	* *	0	* *
Idaho	10,516	78.9 (3)	4,031	27.8 (13)	4,421	15.5 (17)	2,360	30.8 (1)	3,254	20.9 (1)	0	* *	0	* *
Illinois	84,778	61.2 (18)	64,077	18.2 (30)	45,227	9.2 (30)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7D
2006 Pediatric Nutrition Surveillance National (1)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4,5)

Contributor	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Indiana	41,736	60.0 (22)	23,955	19.2 (28)	25,281	9.2 (30)	16,827	6.8 (12)	21,580	2.6 (12)	55,880	62.5 (13)	139,217	30.7 (17)
Iowa	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *
Kansas	8,666	66.0 (15)	5,708	22.2 (24)	3,469	11.8 (27)	0	* *	0	* *	20,943	76.7 (5)	51,130	19.0 (13)
Kentucky	0	* *	0	* *	0	* *	0	* *	0	* *	59,198	98.8 (1)	135,754	32.7 (18)
Maryland	32,588	55.5 (27)	12,412	39.6 (2)	11,282	5.7 (36)	0	* *	0	* *	40,875	75.4 (9)	94,643	7.4 (3)
Massachusetts	30,232	70.1 (9)	11,800	26.2 (14)	16,170	13.0 (24)	0	* *	0	* *	0	* *	0	* *
Michigan	58,676	51.2 (33)	41,890	15.8 (33)	36,303	12.9 (25)	0	* *	0	* *	0	* *	0	* *
Minnesota	32,520	71.0 (8)	12,104	32.6 (7)	23,311	19.2 (9)	6,512	14.4 (5)	12,011	10.1 (3)	0	* *	0	* *
Missouri	43,117	51.6 (32)	12,680	21.9 (25)	24,762	18.8 (11)	0	* *	0	* *	0	* *	0	* *
Montana	5,004	74.5 (5)	1,716	31.5 (10)	2,154	18.9 (10)	0	* *	0	* *	0	* *	0	* *
Navajo Nation	3,071	75.4 (4)	1,323	30.8 (11)	2,300	24.7 (6)	0	* *	0	* *	0	* *	0	* *
Nebraska	11,546	67.8 (12)	7,732	24.4 (18)	6,499	16.2 (15)	3,824	16.2 (4)	4,583	8.4 (5)	20,215	81.9 (2)	45,086	29.4 (16)
Nevada	14,383	54.2 (30)	9,100	23.4 (21)	6,134	14.6 (20)	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7D
2006 Pediatric Nutrition Surveillance National (1)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4,5)

Contributor	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
New Hampshire	4,579	64.6 (16)	1,758	22.8 (23)	2,317	14.7 (19)	995	21.7 (2)	1,605	9.2 (4)	6,069	80.3 (3)	13,730	41.0 (19)
New Jersey	43,359	58.9 (23)	34,042	33.7 (6)	23,691	26.0 (5)	15,008	8.5 (10)	18,648	8.0 (6)	63,690	37.5 (14)	156,959	10.3 (7)
New Mexico	16,194	71.5 (7)	10,638	23.8 (20)	10,654	20.7 (8)	5,503	20.2 (3)	8,053	13.0 (2)	0	* *	0	* *
New York	119,038	69.0 (11)	41,796	39.3 (3)	82,878	24.0 (7)	16,462	10.2 (9)	27,456	3.7 (10)	220,287	76.3 (7)	464,077	2.9 (1)
North Carolina	58,967	57.1 (26)	20,152	19.6 (27)	29,092	14.9 (18)	0	* *	0	* *	0	* *	0	* *
Ohio Child Hlth	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *	27,757	8.5 (5)
Ohio-WIC	73,322	44.4 (35)	45,624	16.6 (32)	33,386	10.5 (29)	0	* *	0	* *	0	* *	0	* *
Oregon	21,813	90.4 (1)	13,044	44.5 (1)	13,191	27.6 (2)	0	* *	0	* *	0	* *	0	* *
Pennsylvania	66,420	42.6 (36)	22,817	20.8 (26)	38,959	11.9 (26)	0	* *	0	* *	0	* *	0	* *
Puerto Rico	39,205	54.5 (28)	22,897	13.9 (35)	24,316	5.1 (38)	0	* *	0	* *	0	* *	0	* *
Rhode Island	4,486	54.3 (29)	2,365	17.3 (31)	3,271	9.2 (30)	1,094	14.0 (6)	1,886	7.5 (8)	8,530	70.9 (10)	17,761	17.2 (11)
Rosebud Sioux-SD	117	60.7 (20)	49	* *	104	17.3 (14)	15	* *	35	* *	8	* *	1,322	17.9 (12)
South Carolina	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) TV Viewing and Smoking in Household data are not reported by all contributors.
- (3) Analyses based on one record per child.
- (4) Excludes records with unknown data or errors.
- (5) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
- (7) Children ages 2 years and older included in the analyses.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7D
2006 Pediatric Nutrition Surveillance National (1)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4,5)

Contributor	Breastfeeding (5)		Exclusive Breastfeeding (6)		≤2 Hours TV Viewing/Day (7)		Smoking in Household	
	Number (Rank)	% Ever Breastfed (Rank)	Number (Rank)	% Breastfed At Least 6 Months (Rank)	Number (Rank)	% Breastfed At Least 3 Months (Rank)	Number (Rank)	% (Rank)
South Dakota	0	* *	0	* *	0	* *	0	* *
Standing Rock-ND	79	* *	24	* *	54	* *	23	* *
Tennessee	46,777	40.7 (38)	14,705	15.6 (34)	25,066	8.5 (33)	0	* *
Texas	230,742	67.6 (13)	81,249	37.8 (5)	140,429	34.6 (1)	0	* *
Vermont	2,345	69.9 (10)	1,639	31.8 (8)	1,691	14.3 (22)	0	* *
Virginia	31,369	58.3 (24)	11,977	23.1 (22)	23,045	16.0 (16)	5,347	6.1 (13)
Washington	42,594	83.1 (2)	32,747	38.8 (4)	22,639	26.8 (4)	0	* *
West Virginia	13,389	42.1 (37)	9,146	11.8 (36)	5,744	7.0 (35)	9,101	3.6 (14)
Wisconsin	31,071	62.6 (17)	12,726	26.0 (15)	17,467	13.2 (23)	8,396	13.7 (7)
Nation	1,577,832	60.1	790,341	25.2	875,205	18.1	104,492	10.3
							143,261	5.6
							584,732	73.3
							2,855,911	10.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 8D
2006 Pediatric Nutrition Surveillance (1)
National
Growth and Anemia Indicators by Race/Ethnicity or Age
Children Aged < 5 Years (2)

Growth and Anemia Indicators by Race/Ethnicity or Age (3)												
	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Number	Low %	High %	Number	Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		<2500 g	>4000 g		% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th		
Race/Ethnicity												
White, Not Hispanic	627,003	8.9	7.4	2,307,899	6.5	4.7	11.7	943,306	15.9	12.3	1,450,549 11.1	
Black, Not Hispanic	389,926	13.5	4.2	1,335,393	7.1	6.1	12.1	535,781	14.4	11.9	825,012 21.0	
Hispanic	692,875	7.2	7.1	2,926,333	5.6	3.8	16.3	1,261,632	17.8	18.0	1,922,871 13.5	
American Indian/Alaskan Native	18,065	8.1	9.0	73,834	5.0	3.3	18.3	33,488	20.1	19.4	49,163 11.5	
Asian/Pacific Islander	45,027	8.3	4.8	179,402	6.4	6.2	11.0	72,109	14.0	12.4	101,446 12.5	
Multiple Races	33,697	9.3	7.0	111,120	5.9	4.2	12.7	41,358	16.1	13.0	64,615 15.0	
All Other	36,377	7.1	7.9	258,062	9.9	6.9	14.8	57,538	16.4	15.6	107,935 11.9	
Total	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591 14.0	
Age												
0 - 11 Months	-	-	-	2,666,703	8.8	6.5	10.9	-	*	*	456,731 16.9	
12 - 23 Months	-	-	-	1,580,128	6.3	3.2	17.0	-	*	*	1,523,914 16.2	
24 - 35 Months	-	-	-	1,108,832	4.4	4.0	13.2	1,108,832	15.8	13.2	976,762 15.3	
36 - 47 Months	-	-	-	955,784	4.3	3.9	15.3	955,784	16.5	15.3	861,378 11.5	
48 - 59 Months	-	-	-	880,596	3.8	3.6	16.4	880,596	17.1	16.4	702,806 8.9	
Total	-	-	-	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591 14.0	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 9D
2006 Pediatric Nutrition Surveillance (1)
National

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (2)
Children Aged < 5 Years (3)

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (4)										
	Breastfeeding (5)						<2 Hours TV/Video Viewing/Day (6)		Smoking in Household	
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	559,606	56.6	282,443	20.0	293,598	13.3	237,537	81.1	730,678	24.0
Black, Not Hispanic	362,220	47.3	165,104	18.2	177,375	11.9	125,081	66.1	412,263	10.5
Hispanic	525,929	72.9	275,991	35.6	336,971	26.5	178,043	68.1	1,357,231	4.3
American Indian/Alaskan Native	16,182	65.8	7,776	29.8	9,852	20.9	7,166	73.9	22,939	11.6
Asian/Pacific Islander	37,151	59.7	17,280	29.7	20,982	19.2	24,006	72.3	106,360	5.8
Multiple Races	32,199	58.5	16,430	19.4	16,338	12.0	12,719	73.8	37,836	21.3
All Other	44,545	55.7	25,317	14.9	20,089	6.4	180	81.1	188,609	4.8
Total	1,577,832	60.1	790,341	25.2	875,205	18.1	584,732	73.3	2,855,916	10.6
Age										
0 - 11 Months	-	-	-	-	-	-	0	*	1,142,908	9.9
12 - 23 Months	-	-	-	-	-	-	0	*	621,740	10.7
24 - 35 Months	-	-	-	-	-	-	242,391	76.3	404,188	11.6
36 - 47 Months	-	-	-	-	-	-	180,629	71.6	351,096	11.4
48 - 59 Months	-	-	-	-	-	-	161,712	70.7	335,984	10.9
Total	1,577,832	60.1	790,341	25.2	875,205	18.1	584,732	73.3	2,855,916	10.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 10D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years (2)

Summary of Trends in Racial and Ethnic Distribution								
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
2006	7,599,318	31.5	18.5	41.1	1.0	2.5	1.5	3.9
2005	7,117,562	32.3	18.5	37.7	1.0	2.5	1.0	7.0
2004	6,930,341	35.5	20.3	37.1	1.2	2.9	0.1	2.9
2003	6,358,666	35.5	19.6	37.5	1.1	3.2	0.0	3.0
2002	5,518,837	39.1	21.3	32.0	1.2	3.2	0.0	3.2
2001	4,943,250	42.0	22.3	28.5	1.4	2.9	0.0	2.8
2000	5,019,497	42.3	21.8	28.3	1.5	3.0	0.0	3.2
1999	5,542,296	41.6	23.6	27.7	1.0	3.0	0.0	3.1
1998	5,697,317	42.4	23.6	26.8	1.1	3.1	0.0	3.0
1997	5,703,221	43.4	23.5	25.4	1.4	2.8	0.0	3.4
1996	6,024,305	43.3	23.6	26.2	1.5	2.7	0.0	2.7
1995	5,984,296	43.2	24.4	25.8	1.5	2.7	0.0	2.5
1994	5,711,154	42.9	24.4	26.1	1.5	2.8	0.0	2.3
1993	4,937,857	41.5	24.2	27.6	1.5	2.8	0.0	2.5
1992	4,543,683	40.8	25.0	27.3	1.5	2.7	0.0	2.7
1991	4,319,881	42.3	26.1	25.1	1.3	2.6	0.0	2.5
1990	3,742,602	40.9	27.0	25.4	1.6	2.7	0.0	2.5
1989	3,314,772	42.2	28.0	23.3	1.7	2.7	0.0	2.1
1988	2,036,649	50.3	30.4	15.0	2.4	1.6	0.0	0.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 10D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years (2)

Summary of Trends in Racial and Ethnic Distribution								
Year	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1987	1,738,567	48.3	31.8	16.1	2.3	1.3	0.0	0.2
1986	1,586,020	48.1	31.0	17.0	2.3	1.4	0.0	0.2
1985	1,334,323	47.0	29.9	18.8	2.6	1.5	0.0	0.2
1984	1,054,961	51.2	31.1	13.1	2.7	1.7	0.0	0.3
1983	866,238	53.5	30.5	11.6	2.7	1.4	0.0	0.3
1982	662,762	53.5	31.6	11.6	1.5	1.3	0.0	0.4
1981	528,895	54.2	30.7	11.3	1.8	1.3	0.0	0.6
1980	594,245	51.1	29.8	11.9	5.1	1.0	0.0	1.0
1979	324,360	54.0	26.7	12.0	5.7	0.7	0.0	0.8
1978	239,882	56.2	26.0	9.5	7.2	0.5	0.0	0.6
1977	198,928	55.9	25.9	8.9	8.0	0.4	0.0	0.9
1976	150,400	53.2	20.7	11.7	12.6	0.3	0.0	1.4
1975	113,146	52.5	20.8	10.9	13.2	0.2	0.0	2.4
1974	52,315	59.0	18.5	14.3	5.7	0.2	0.0	2.4
1973	12,075	19.0	80.6	0.2	0.1	0.0	0.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Summary of Trends in Age Distribution (3)				
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %
2006	7,599,318	37.6	21.8	40.6
2005	7,117,562	35.3	22.0	42.6
2004	6,930,341	35.4	22.2	42.5
2003	6,358,666	35.8	21.9	42.3
2002	5,518,837	36.5	21.7	41.8
2001	4,943,250	37.8	21.4	40.8
2000	5,019,497	37.7	21.2	41.1
1999	5,542,296	36.9	21.3	41.8
1998	5,697,317	36.1	21.2	42.7
1997	5,703,221	35.1	20.9	44.0
1996	6,024,305	35.3	21.2	43.6
1995	5,984,296	35.8	21.6	42.6
1994	5,711,154	37.0	21.7	41.2
1993	4,937,857	37.3	22.7	40.1
1992	4,543,683	39.6	23.2	37.3
1991	4,319,881	42.9	22.3	34.8
1990	3,742,602	43.5	22.0	34.6
1989	3,314,772	42.5	22.2	35.3

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Records rejected if date of birth and date of visit unknown.
(4) Excludes records with unknown data or errors.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Summary of Trends in Age Distribution (3)				
Year	Number Included in Analysis (4)	0-11 Months %	12-23 Months %	24-59 Months %
1988	2,036,649	42.8	22.3	34.9
1987	1,738,567	43.4	22.1	34.4
1986	1,586,020	44.0	21.1	34.9
1985	1,334,323	42.5	20.8	36.7
1984	1,054,961	42.6	21.3	36.1
1983	866,238	44.8	21.6	33.6
1982	662,762	43.7	21.1	35.2
1981	528,895	42.6	21.7	35.7
1980	594,245	37.8	22.4	39.7
1979	324,360	36.5	22.6	41.0
1978	239,882	37.0	22.4	40.6
1977	198,928	36.2	21.8	42.0
1976	150,400	34.0	22.8	43.2
1975	113,146	37.9	23.2	38.9
1974	52,315	40.7	21.9	37.4
1973	12,075	16.4	19.3	64.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Records rejected if date of birth and date of visit unknown.

(4) Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Number	%	
	Number	% <2500 g	% >4000 g									Number
2006	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591	14.0
2005	1,641,264	9.4	6.6	6,709,486	6.4	4.8	13.9	2,869,075	16.1	14.7	4,381,087	13.6
2004	1,630,186	9.3	6.8	6,582,905	6.4	4.7	14.1	2,814,475	16.1	14.8	4,364,901	13.5
2003	1,505,016	9.0	7.2	6,003,860	6.4	5.0	13.9	2,576,591	15.8	14.7	3,600,353	13.4
2002	1,254,683	9.0	7.6	5,060,321	6.3	5.5	13.5	2,139,100	15.4	14.3	3,421,187	13.0
2001	1,217,626	9.0	8.0	4,390,867	6.3	5.4	12.9	1,814,873	15.1	13.1	3,116,606	13.4
2000	1,235,686	8.9	8.3	4,617,585	6.0	5.4	12.8	1,932,131	15.2	13.2	3,217,568	13.7
1999	1,301,847	8.9	8.4	5,149,796	5.9	5.7	12.3	2,187,151	15.0	12.7	3,645,746	14.5
1998	1,307,687	8.9	8.5	5,365,056	6.0	5.9	12.0	2,320,924	14.9	12.2	3,864,299	14.5
1997	1,199,049	8.8	8.5	5,389,269	6.1	5.9	11.7	2,396,884	14.6	11.6	3,925,299	14.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Number	%	
	Number	% <2500 g	% >4000 g									Number
1996	1,277,355	8.7	8.7	5,624,304	6.2	6.0	11.4	2,479,125	14.4	11.4	4,158,251	15.3
1995	1,289,474	9.0	8.6	5,497,483	6.6	6.0	11.2	2,372,868	14.1	11.0	4,125,108	15.9
1994	1,270,449	9.2	8.5	5,192,014	6.8	6.2	11.2	2,168,677	13.9	10.9	3,855,156	15.8
1993	1,061,208	9.2	8.4	4,566,496	6.8	6.7	11.0	1,857,797	13.6	10.7	3,368,409	16.1
1992	962,937	9.3	8.4	4,189,029	6.9	6.9	10.9	1,585,600	13.5	10.6	3,016,330	15.9
1991	1,020,919	9.4	8.3	3,977,097	7.1	7.3	10.5	1,403,114	13.0	10.0	2,748,979	16.3
1990	958,828	9.2	8.2	3,329,394	7.4	7.5	10.6	1,156,421	13.1	10.0	2,362,182	16.3
1989	826,736	9.1	8.1	3,106,714	7.4	7.4	10.4	1,115,794	12.9	9.6	2,111,468	15.6
1988	532,664	9.5	7.8	1,903,637	7.8	7.8	9.1	676,741	12.1	8.4	1,380,353	14.8
1987	444,229	9.9	7.5	1,668,076	8.2	8.1	9.2	585,297	11.9	8.2	1,217,664	14.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under-weight (6)	Over-weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1986	367,997	9.9	7.6	1,512,514	8.5	8.0	9.4	538,821	12.0	8.4	1,127,586	15.0
1985	313,159	10.0	7.4	1,262,066	8.4	8.3	9.3	473,634	11.9	8.2	966,479	15.9
1984	249,655	10.0	7.6	989,258	8.6	7.9	8.8	368,151	12.0	7.6	770,212	16.6
1983	217,314	10.3	7.6	808,045	9.4	8.0	8.8	282,979	12.1	7.7	619,567	17.9
1982	168,673	10.2	7.8	617,201	9.8	7.7	9.1	227,461	12.7	8.1	466,815	18.1
1981	119,475	10.5	7.9	493,366	9.7	7.5	9.8	184,223	13.0	8.5	371,131	19.5
1980	124,800	10.0	8.5	518,312	8.8	7.2	10.0	214,369	13.0	8.6	432,280	18.1
1979	67,299	10.1	8.2	270,921	8.5	6.8	10.0	113,329	12.8	8.3	240,211	16.8
1978	48,187	10.0	8.3	207,748	8.7	7.3	9.8	84,956	12.6	8.2	177,846	18.9
1977	40,767	10.2	8.6	170,103	9.0	7.7	10.7	72,267	12.8	8.4	146,205	18.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators (3)												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1976	26,045	9.6	8.7	121,153	9.3	7.1	12.0	53,521	13.4	9.6	114,145	16.4
1975	20,728	9.7	8.4	85,611	10.5	7.8	12.5	33,864	13.5	10.3	81,821	19.3
1974	10,309	9.8	9.1	34,099	11.7	8.8	12.7	12,897	14.2	10.5	36,301	22.4
1973	195	9.7	11.8	11,078	8.7	8.4	10.2	7,375	10.2	6.6	10,884	31.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Breastfeeding
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding (3)						
Year	Breastfeeding (4)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
2006	1,577,832	60.1	790,341	25.2	875,205	18.1
2005	1,515,954	58.5	751,259	24.3	869,342	17.7
2004	1,466,352	56.1	730,949	23.4	816,328	17.5
2003	1,244,877	55.1	606,003	23.4	701,977	17.3
2002	1,014,091	52.5	546,600	20.7	582,948	12.3
2001	553,594	50.1	311,294	21.5	310,791	13.9
2000	905,154	48.0	504,089	19.7	478,170	12.3
1999	864,433	47.8	495,446	18.6	452,736	11.7
1998	877,246	47.8	490,802	18.1	475,699	11.8
1997	750,078	45.1	440,557	16.8	416,287	11.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Breastfeeding
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding (3)						
Year	Breastfeeding (4)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
1996	796,109	43.7	479,281	15.6	461,059	10.1
1995	717,537	41.6	409,439	16.0	401,859	10.7
1994	692,196	39.2	379,703	14.7	363,662	10.1
1993	479,000	38.9	266,051	15.0	271,481	11.6
1992	374,892	36.3	243,882	15.1	230,718	10.7
1991	511,841	32.5	258,025	13.7	241,238	10.6
1990	403,472	38.1	204,682	13.1	175,847	9.8
1989	347,322	37.9	185,009	13.4	160,759	10.3
1988	324,476	37.0	179,168	12.8	144,450	9.7
1987	266,327	39.4	157,403	12.7	123,455	10.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Breastfeeding
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding (3)						
Year	Breastfeeding (4)					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
1986	218,391	42.6	144,971	14.1	102,730	10.8
1985	173,368	46.0	116,531	14.3	78,866	10.8
1984	137,267	42.7	89,165	12.1	61,384	8.1
1983	34,661	41.3	23,107	14.5	15,338	10.5
1982	9,242	35.5	5,759	17.1	4,282	9.9
1981	1,274	32.4	1,263	8.2	1,096	7.1
1980	1,081	28.0	1,053	9.0	842	4.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 14D
2006 Pediatric Nutrition Surveillance
National
Children Aged < 5 Years

Page: 1
Run Date: 8 / 27 / 07

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 15D
2006 Pediatric Nutrition Surveillance
National
Children Aged < 5 Years

Page: 1
Run Date: 8 / 27 / 07

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 16D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Growth Indicators by Race/Ethnicity and Age ⁽³⁾									
	Short Stature (4)		Underweight (5)		Overweight (5)		≥ 2 Yrs Overweight (5)		
	Number	% <5th	Number	% <5th	Number	% ≥95th	Number	% 85th - <95th	% ≥95th
White, Not Hispanic									
0 - 11 Months	869,286	8.8	869,286	6.6	869,286	9.2	0	0.0	0.0
12 - 23 Months	495,307	6.3	495,307	3.0	495,307	15.0	0	0.0	0.0
24 - 59 Months	943,306	4.5	943,306	3.7	943,306	12.3	943,306	15.9	12.3
Total	2,307,899	6.5	2,307,899	4.7	2,307,899	11.7	943,306	15.9	12.3
Black, Not Hispanic									
0 - 11 Months	503,374	11.2	503,374	8.0	503,374	10.3	0	0.0	0.0
12 - 23 Months	296,238	6.6	296,238	4.2	296,238	15.6	0	0.0	0.0
24 - 59 Months	535,781	3.4	535,781	5.4	535,781	11.9	535,781	14.4	11.9
Total	1,335,393	7.1	1,335,393	6.1	1,335,393	12.1	535,781	14.4	11.9
Hispanic									
0 - 11 Months	1,012,455	7.2	1,012,455	5.4	1,012,455	12.3	0	0.0	0.0
12 - 23 Months	652,246	6.0	652,246	2.5	652,246	19.3	0	0.0	0.0
24 - 59 Months	1,261,632	4.2	1,261,632	3.2	1,261,632	18.0	1,261,632	17.8	18.0
Total	2,926,333	5.6	2,926,333	3.8	2,926,333	16.3	1,261,632	17.8	18.0
American Indian/Alaskan Native									
0 - 11 Months	23,624	7.6	23,624	5.5	23,624	12.6	0	0.0	0.0
12 - 23 Months	16,722	5.0	16,722	2.0	16,722	24.0	0	0.0	0.0
24 - 59 Months	33,488	3.2	33,488	2.4	33,488	19.4	33,488	20.1	19.4
Total	73,834	5.0	73,834	3.3	73,834	18.3	33,488	20.1	19.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 16D
2006 Pediatric Nutrition Surveillance ⁽¹⁾
National
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Growth Indicators by Race/Ethnicity and Age ⁽³⁾									
	Short Stature (4)		Underweight (5)		Overweight (5)		≥ 2 Yrs Overweight (5)		
	Number	% <5th	Number	% <5th	Number	% ≥95th	Number	% 85th - <95th	% ≥95th
Asian/Pacific Islander									
0 - 11 Months	67,957	7.1	67,957	7.2	67,957	9.3	0	0.0	0.0
12 - 23 Months	39,336	6.9	39,336	5.9	39,336	11.6	0	0.0	0.0
24 - 59 Months	72,109	5.6	72,109	5.5	72,109	12.4	72,109	14.0	12.4
Total	179,402	6.4	179,402	6.2	179,402	11.0	72,109	14.0	12.4
Multiple Races									
0 - 11 Months	45,312	8.6	45,312	5.4	45,312	10.2	0	0.0	0.0
12 - 23 Months	24,450	5.2	24,450	2.6	24,450	16.6	0	0.0	0.0
24 - 59 Months	41,358	3.3	41,358	3.7	41,358	13.0	41,358	16.1	13.0
Total	111,120	5.9	111,120	4.2	111,120	12.7	41,358	16.1	13.0
All Other/Unknown									
0 - 11 Months	144,695	12.9	144,695	8.0	144,695	13.5	0	0.0	0.0
12 - 23 Months	55,829	8.0	55,829	5.4	55,829	17.3	0	0.0	0.0
24 - 59 Months	57,538	4.3	57,538	5.4	57,538	15.6	57,538	16.4	15.6
Total	258,062	9.9	258,062	6.9	258,062	14.8	57,538	16.4	15.6
Total All Race/Ethnic Groups									
0 - 11 Months	2,666,703	8.8	2,666,703	6.5	2,666,703	10.9	0	0.0	0.0
12 - 23 Months	1,580,128	6.3	1,580,128	3.2	1,580,128	17.0	0	0.0	0.0
24 - 59 Months	2,945,212	4.2	2,945,212	3.9	2,945,212	14.8	2,945,212	16.4	14.8
Total	7,192,043	6.4	7,192,043	4.7	7,192,043	13.8	2,945,212	16.4	14.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17D
2006 Pediatric Nutrition Surveillance (1)
National
Anemia Indicators by Race/Ethnicity and Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 8 / 27 / 07

Anemia Indicators by Race/Ethnicity and Age (3, 4)					
	Anemia Low Hb/Hct %			Anemia Low Hb/Hct %	
	Number		Number		
White, Not Hispanic			Asian/Pacific Islander		
6 - 11 Months	156,951	15.1	6 - 11 Months	11,169	13.1
12 - 17 Months	295,376	14.2	12 - 17 Months	20,674	12.8
18 - 23 Months	202,254	11.3	18 - 23 Months	13,279	12.5
24 - 35 Months	314,876	11.5	24 - 35 Months	21,942	14.9
36 - 59 Months	481,092	7.5	36 - 59 Months	34,382	10.7
Total	1,450,549	11.1	Total	101,446	12.5
Black, Not Hispanic			Multiple Races		
6 - 11 Months	77,420	21.6	6 - 11 Months	6,843	17.1
12 - 17 Months	179,143	23.1	12 - 17 Months	14,668	18.1
18 - 23 Months	111,772	22.7	18 - 23 Months	9,571	15.1
24 - 35 Months	180,474	23.1	24 - 35 Months	13,920	15.6
36 - 59 Months	276,203	17.4	36 - 59 Months	19,613	11.5
Total	825,012	21.0	Total	64,615	15.0
Hispanic			All Other/Unknown		
6 - 11 Months	171,603	17.4	6 - 11 Months	28,459	12.5
12 - 17 Months	377,410	17.1	12 - 17 Months	29,615	10.8
18 - 23 Months	245,423	14.6	18 - 23 Months	8,683	11.6
24 - 35 Months	419,420	14.9	24 - 35 Months	15,067	14.7
36 - 59 Months	709,015	9.5	36 - 59 Months	26,111	10.8
Total	1,922,871	13.5	Total	107,935	11.9
American Indian/Alaskan Native			Total All Race/Ethnic Groups		
6 - 11 Months	4,286	15.2	6 - 11 Months	456,731	16.9
12 - 17 Months	9,168	15.3	12 - 17 Months	926,054	17.0
18 - 23 Months	6,878	13.9	18 - 23 Months	597,860	14.9
24 - 35 Months	11,063	13.2	24 - 35 Months	976,762	15.3
36 - 59 Months	17,768	6.8	36 - 59 Months	1,564,184	10.3
Total	49,163	11.5	Total	4,521,591	14.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
White, Not Hispanic	2006	627,003	8.9	7.4	2,307,899	6.5	4.7	11.7	943,306	15.9	12.3	1,450,549	11.1
	2005	574,328	9.1	7.5	2,205,721	6.6	4.6	12.0	928,533	15.7	12.4	1,434,886	10.8
	2004	629,412	8.8	7.8	2,367,268	6.6	4.5	12.2	994,087	15.8	12.6	1,549,047	10.8
	2003	567,095	8.6	8.3	2,154,320	6.5	4.8	11.9	920,156	15.5	12.2	1,337,637	10.5
	2002	522,252	8.4	8.8	1,980,732	6.3	5.0	11.5	850,084	15.0	11.8	1,374,440	10.1
	2001	536,570	8.4	9.2	1,845,533	6.3	5.2	11.3	774,389	14.8	11.5	1,347,726	10.4
	2000	548,613	8.3	9.6	1,977,676	6.0	5.3	11.1	832,279	14.9	11.3	1,393,526	10.6
	1999	573,074	8.3	9.8	2,177,307	6.0	5.4	10.4	920,094	14.6	10.6	1,537,766	11.3
	1998	592,140	8.1	10.0	2,314,653	6.1	5.7	10.0	997,828	14.4	10.0	1,661,249	11.1
	1997	560,695	8.1	10.0	2,378,409	6.3	5.6	9.9	1,049,613	14.2	9.6	1,725,769	11.4
Black, Not Hispanic	2006	389,926	13.5	4.2	1,335,393	7.1	6.1	12.1	535,781	14.4	11.9	825,012	21.0
	2005	330,923	13.6	4.4	1,247,201	6.9	6.1	12.4	527,365	14.3	12.0	793,328	20.0
	2004	356,415	13.1	4.5	1,334,648	6.9	6.0	12.8	563,849	14.4	12.2	874,513	19.4
	2003	318,381	12.9	4.8	1,173,486	7.1	6.4	12.4	494,997	14.1	12.1	724,232	19.1
	2002	286,273	12.6	4.9	1,094,418	7.0	6.7	12.0	453,415	13.7	11.9	742,716	19.1
	2001	300,376	12.7	5.0	1,014,897	7.5	6.7	11.8	408,557	13.6	11.0	707,241	19.6
	2000	299,774	12.6	5.2	1,027,120	7.2	6.7	11.7	419,361	13.7	11.1	713,877	20.0
	1999	332,583	12.6	5.3	1,228,248	6.9	7.2	11.3	516,002	13.6	10.8	876,387	21.2
	1998	332,792	12.6	5.3	1,277,982	6.9	7.5	11.1	555,698	13.5	10.6	938,727	21.0
	1997	284,641	12.5	5.3	1,280,130	6.9	7.3	10.9	592,353	13.4	10.2	956,175	20.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
Hispanic	2006	692,875	7.2	7.1	2,926,333	5.6	3.8	16.3	1,261,632	17.8	18.0	1,922,871	13.5
	2005	548,820	7.4	7.1	2,519,801	5.9	4.0	16.6	1,134,627	17.5	18.0	1,735,800	13.2
	2004	544,571	7.7	7.1	2,423,347	6.0	4.0	16.8	1,076,000	17.4	18.2	1,684,460	13.1
	2003	522,353	7.3	7.5	2,238,214	6.0	4.3	16.8	991,223	17.0	18.4	1,301,313	13.3
	2002	360,889	7.1	8.0	1,602,567	6.0	5.0	17.0	687,055	17.0	19.0	1,081,779	12.5
	2001	304,986	6.7	8.7	1,223,159	5.5	4.5	16.1	509,679	16.4	17.0	871,516	13.0
	2000	307,503	6.7	9.0	1,275,430	5.2	4.5	16.2	541,044	16.6	17.2	903,791	13.3
	1999	318,342	6.7	9.1	1,398,874	5.0	4.8	15.8	607,461	16.6	16.9	1,015,597	13.8
	1998	307,317	6.7	8.9	1,415,043	5.0	4.8	15.8	613,072	16.8	16.7	1,028,705	14.3
	1997	267,469	6.8	8.8	1,340,845	5.2	4.9	15.3	590,482	16.1	16.0	984,513	14.3
American Indian/Alaskan Native	2006	18,065	8.1	9.0	73,834	5.0	3.3	18.3	33,488	20.1	19.4	49,163	11.5
	2005	16,080	7.9	9.2	69,413	5.1	3.1	18.4	32,308	20.0	19.0	49,165	10.7
	2004	18,468	7.9	9.8	76,679	5.0	3.3	18.7	35,328	19.9	19.0	54,388	11.5
	2003	16,191	7.4	10.2	68,766	5.1	3.6	17.4	31,798	18.9	17.7	48,437	10.9
	2002	14,420	6.8	11.0	58,382	5.1	3.4	17.2	28,324	18.8	17.5	45,762	10.0
	2001	15,339	7.3	10.9	59,571	5.2	3.5	16.7	28,917	18.9	17.1	49,255	9.9
	2000	16,066	6.7	11.7	67,463	5.1	3.2	17.5	32,159	19.9	17.4	50,399	11.0
	1999	12,807	6.8	12.4	53,043	5.0	4.2	16.9	24,645	19.2	17.2	38,719	12.9
	1998	13,719	7.4	12.3	59,983	4.8	4.2	16.5	28,832	19.3	16.5	44,741	13.4
	1997	17,105	7.0	12.2	79,184	4.9	4.3	15.5	39,425	19.2	15.4	60,037	12.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
Asian/Pacific Islander	2006	45,027	8.3	4.8	179,402	6.4	6.2	11.0	72,109	14.0	12.4	101,446	12.5
	2005	39,522	8.5	5.1	159,167	6.6	6.6	11.1	65,934	14.1	12.3	100,090	12.4
	2004	43,519	8.4	5.7	188,721	6.7	6.0	11.7	75,018	13.9	12.5	106,595	12.7
	2003	44,761	8.3	5.9	193,549	6.0	6.1	11.9	73,816	14.1	13.0	101,017	12.6
	2002	36,052	8.0	6.2	158,642	6.4	6.2	12.1	61,152	13.8	13.4	94,638	12.4
	2001	33,340	8.3	6.2	122,338	7.0	5.5	12.4	48,926	14.5	13.4	76,669	12.8
	2000	34,139	8.0	6.7	137,405	6.7	5.6	12.6	59,924	14.5	13.8	83,969	15.1
	1999	34,690	7.7	6.8	154,695	6.6	6.6	11.7	69,312	14.6	13.3	98,523	13.8
	1998	33,174	7.9	6.6	164,874	6.6	6.5	12.1	78,491	15.0	13.9	113,199	15.2
	1997	30,906	7.9	6.0	153,123	7.1	6.9	11.2	69,465	14.3	12.9	105,558	12.9
Multiple Races	2006	33,697	9.3	7.0	111,120	5.9	4.2	12.7	41,358	16.1	13.0	64,615	15.0
	2005	21,536	9.6	7.0	55,341	6.5	4.3	12.7	20,450	15.6	13.0	38,478	14.0
	2004	2,185	8.8	6.5	6,483	6.2	3.5	13.9	2,359	15.9	15.1	3,869	17.3
	2003	0	*	*	0	*	*	*	0	*	*	0	*
	2002	0	*	*	0	*	*	*	0	*	*	0	*
	2001	0	*	*	0	*	*	*	0	*	*	0	*
	2000	0	*	*	0	*	*	*	0	*	*	0	*
	1999	0	*	*	0	*	*	*	0	*	*	0	*
	1998	0	*	*	0	*	*	*	0	*	*	0	*
	1997	0	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18D
2006 Pediatric Nutrition Surveillance (1)
National

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
		Low		High	Short Stature (5)		Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)				
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th		
All Other/Unknown	2006	36,377	7.1	7.9	258,062	9.9	6.9	14.8	57,538	16.4	15.6	107,935	11.9
	2005	100,362	9.2	6.7	400,070	7.5	5.9	13.0	141,209	15.7	14.5	208,640	12.9
	2004	35,616	8.1	8.1	185,759	5.4	6.6	13.1	67,834	15.7	15.2	92,029	14.0
	2003	36,235	8.1	8.0	175,525	5.4	6.9	13.0	64,601	15.3	14.9	87,717	13.6
	2002	34,797	8.2	8.2	165,580	5.5	7.1	12.9	59,070	15.0	14.7	81,852	13.7
	2001	27,015	8.1	8.4	125,369	5.7	6.4	13.4	44,405	15.5	14.2	64,199	14.5
	2000	29,591	8.5	8.4	132,491	5.7	6.5	13.3	47,364	15.2	14.6	72,006	14.0
	1999	30,351	8.2	8.9	137,629	5.5	6.6	12.6	49,637	15.3	13.8	78,754	14.6
	1998	28,545	7.8	9.2	132,521	5.5	6.5	12.6	47,003	15.2	13.7	77,678	13.9
	1997	38,233	7.7	8.7	157,578	5.8	6.4	13.1	55,546	15.5	14.5	93,247	14.4
Total All Race/Ethnic Groups	2006	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591	14.0
	2005	1,631,571	9.4	6.6	6,656,714	6.4	4.7	13.9	2,850,426	16.1	14.7	4,360,387	13.6
	2004	1,630,186	9.3	6.8	6,582,905	6.4	4.7	14.1	2,814,475	16.1	14.8	4,364,901	13.5
	2003	1,505,016	9.0	7.2	6,003,860	6.4	5.0	13.9	2,576,591	15.8	14.7	3,600,353	13.4
	2002	1,254,683	9.0	7.6	5,060,321	6.3	5.5	13.5	2,139,100	15.4	14.3	3,421,187	13.0
	2001	1,217,626	9.0	8.0	4,390,867	6.3	5.4	12.9	1,814,873	15.1	13.1	3,116,606	13.4
	2000	1,235,686	8.9	8.3	4,617,585	6.0	5.4	12.8	1,932,131	15.2	13.2	3,217,568	13.7
	1999	1,301,847	8.9	8.4	5,149,796	5.9	5.7	12.3	2,187,151	15.0	12.7	3,645,746	14.5
	1998	1,307,687	8.9	8.5	5,365,056	6.0	5.9	12.0	2,320,924	14.9	12.2	3,864,299	14.5
	1997	1,199,049	8.8	8.5	5,389,269	6.1	5.9	11.7	2,396,884	14.6	11.6	3,925,299	14.5

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Infants born during the reporting period included in the analysis.
(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 19D
2006 Pediatric Nutrition Surveillance (1)**

**National
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)**

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
White, Not Hispanic	2006	559,606	56.6	282,443	20.0	293,598	13.3
	2005	535,489	55.4	273,609	19.7	299,095	13.0
	2004	577,914	53.8	295,108	18.8	310,618	12.7
	2003	479,220	52.6	237,240	19.3	255,873	13.5
	2002	427,916	51.0	235,051	18.6	243,022	12.1
	2001	291,524	50.6	168,134	20.1	165,987	13.3
	2000	428,481	48.5	246,974	18.7	228,170	11.9
	1999	428,232	49.0	256,561	18.1	228,248	11.7
	1998	441,284	50.5	253,111	17.8	240,209	11.7
	1997	405,189	47.8	246,439	17.3	227,333	11.7
Black, Not Hispanic	2006	362,220	47.3	165,104	18.2	177,375	11.9
	2005	327,559	46.4	152,600	18.4	174,080	12.0
	2004	354,100	42.6	163,265	16.3	190,170	11.0
	2003	296,353	41.6	140,154	16.3	160,412	10.7
	2002	267,397	40.0	135,737	15.2	148,196	9.0
	2001	144,375	36.1	73,220	14.0	78,159	9.6
	2000	258,475	34.5	133,425	13.3	131,564	8.2
	1999	244,816	35.3	130,327	12.9	123,557	7.9
	1998	242,275	33.5	129,336	12.1	126,687	7.8
	1997	194,656	30.5	106,634	10.2	102,610	6.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 19D
2006 Pediatric Nutrition Surveillance (1)**

**National
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)**

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Hispanic	2006	525,929	72.9	275,991	35.6	336,971	26.5
	2005	455,203	71.1	245,636	33.2	314,397	25.7
	2004	453,498	68.7	232,953	33.8	276,523	27.3
	2003	398,490	68.0	194,066	33.5	246,698	25.9
	2002	254,397	67.5	144,220	28.7	158,724	15.3
	2001	90,392	68.4	54,084	33.8	50,918	20.3
	2000	171,314	65.5	94,827	30.1	92,202	17.9
	1999	155,326	62.8	86,781	27.7	82,517	16.8
	1998	155,982	61.1	85,802	26.9	86,733	17.3
	1997	103,848	59.4	61,994	24.9	63,408	16.7
American Indian/Alaskan Native	2006	16,182	65.8	7,776	29.8	9,852	20.9
	2005	13,272	61.5	6,889	24.7	8,496	17.7
	2004	14,683	62.4	7,644	20.8	8,642	15.9
	2003	11,858	55.7	6,014	22.3	7,576	17.4
	2002	9,141	58.6	5,214	21.5	5,562	12.1
	2001	5,817	52.2	3,395	22.9	3,260	15.2
	2000	10,134	53.2	6,193	23.2	5,892	17.0
	1999	7,583	55.9	4,730	25.3	4,453	19.2
	1998	11,664	56.7	7,204	22.1	7,438	14.9
	1997	14,447	57.7	7,368	22.8	8,809	17.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19D
2006 Pediatric Nutrition Surveillance (1)

National
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
Asian/Pacific Islander	2006	37,151	59.7	17,280	29.7	20,982	19.2
	2005	39,048	59.4	16,941	29.6	21,245	20.0
	2004	43,207	58.8	22,061	23.6	18,164	19.4
	2003	38,507	53.6	19,516	21.8	20,863	13.0
	2002	32,440	51.0	17,442	21.7	17,343	12.2
	2001	10,641	55.3	6,566	24.6	6,190	16.3
	2000	24,294	54.5	15,091	20.9	14,523	12.6
	1999	18,578	50.3	11,122	20.7	10,078	13.0
	1998	18,307	51.8	10,519	20.7	11,207	12.6
	1997	13,848	47.7	7,977	20.3	7,992	13.3
Multiple Races	2006	32,199	58.5	16,430	19.4	16,338	12.0
	2005	21,346	57.2	9,559	19.1	10,693	12.9
	2004	2,117	50.8	811	19.6	1,209	10.9
	2003	0	*	0	*	0	*
	2002	0	*	0	*	0	*
	2001	0	*	0	*	0	*
	2000	0	*	0	*	0	*
	1999	0	*	0	*	0	*
	1998	0	*	0	*	0	*
	1997	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19D
2006 Pediatric Nutrition Surveillance (1)

National
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Breastfeeding Indicators by Race/Ethnicity (3)							
Race/Ethnicity and Year		Breastfeeding (4)					
		Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
All Other/Unknown	2006	44,545	55.7	25,317	14.9	20,089	6.4
	2005	124,037	57.4	46,025	23.6	41,336	15.2
	2004	20,833	62.7	9,107	31.5	11,002	20.1
	2003	20,449	61.2	9,013	31.4	10,555	19.6
	2002	22,800	57.9	8,936	28.8	10,101	18.9
	2001	10,845	61.1	5,895	35.0	6,277	27.4
	2000	12,456	54.6	7,579	30.4	5,819	21.8
	1999	9,898	54.1	5,925	26.4	3,883	17.2
	1998	7,734	57.4	4,830	25.4	3,425	15.9
	1997	18,090	46.3	10,145	17.5	6,135	13.5
Total All Race/Ethnic Groups	2006	1,577,832	60.1	790,341	25.2	875,205	18.1
	2005	1,515,954	58.5	751,259	24.3	869,342	17.7
	2004	1,466,352	56.1	730,949	23.4	816,328	17.5
	2003	1,244,877	55.1	606,003	23.4	701,977	17.3
	2002	1,014,091	52.5	546,600	20.7	582,948	12.3
	2001	553,594	50.1	311,294	21.5	310,791	13.9
	2000	905,154	48.0	504,089	19.7	478,170	12.3
	1999	864,433	47.8	495,446	18.6	452,736	11.7
	1998	877,246	47.8	490,802	18.1	475,699	11.8
	1997	750,078	45.1	440,557	16.8	416,287	11.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 20D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Summary of Trends in Growth Indicators by Age (3)																		
Age and Year *	Height and Weight									Age and Year *	Height and Weight							
	Short Stature (4)		Under-weight (5)	Over-weight (5)		≥ 2 Yrs Overweight (5)					Short Stature (4)		Under-weight (5)	Over-weight (5)		≥ 2 Yrs Overweight (5)		
	Number	%	%	%	Number	%	%	Number			%	%	Number	%	%	Number	%	%
	<5th	<5th	≥95th		85th-<95th	≥95th		<5th	<5th	≥95th		85th-<95th	≥95th		85th-<95th	≥95th		
0 - 11 Months									24 - 35 Months									
2006	2,666,703	8.8	6.5	10.9	0	*	*	2006	1,108,832	4.4	4.0	13.2	1,108,832	15.8	13.2			
2005	2,341,698	9.1	6.7	10.8	0	*	*	2005	1,074,186	4.5	4.2	13.1	1,074,186	15.5	13.1			
2004	2,286,920	9.0	6.5	11.0	0	*	*	2004	1,052,156	4.7	4.2	13.3	1,052,156	15.6	13.3			
2003	2,089,060	8.7	6.8	10.8	0	*	*	2003	969,160	4.6	4.5	13.4	969,160	15.3	13.4			
2002	1,796,771	8.4	7.4	10.4	0	*	*	2002	812,482	4.5	5.0	12.9	812,482	14.7	12.9			
2001	1,608,934	8.5	7.3	10.5	0	*	*	2001	687,768	4.4	4.9	11.9	687,768	14.4	11.9			
2000	1,684,907	8.0	7.6	10.2	0	*	*	2000	719,739	4.3	4.7	11.9	719,739	14.4	11.9			
1999	1,845,669	7.8	8.1	9.8	0	*	*	1999	808,198	4.2	4.9	11.6	808,198	14.2	11.6			
1998	1,882,796	7.9	8.6	9.4	0	*	*	1998	838,169	4.4	5.1	11.2	838,169	14.1	11.2			
1997	1,847,702	8.0	8.3	9.6	0	*	*	1997	852,437	4.6	5.2	10.7	852,437	13.8	10.7			
12 - 23 Months									36 - 47 Months									
2006	1,580,128	6.3	3.2	17.0	0	*	*	2006	955,784	4.3	3.9	15.3	955,784	16.5	15.3			
2005	1,498,713	6.3	3.1	17.3	0	*	*	2005	934,386	4.4	4.1	15.2	934,386	16.2	15.2			
2004	1,481,510	6.4	3.1	17.6	0	*	*	2004	923,436	4.5	4.2	15.3	923,436	16.2	15.3			
2003	1,338,209	6.4	3.3	17.3	0	*	*	2003	839,220	4.5	4.5	15.2	839,220	15.8	15.2			
2002	1,124,450	6.6	3.7	16.8	0	*	*	2002	695,636	4.5	4.8	14.8	695,636	15.5	14.8			
2001	967,060	6.5	3.4	16.6	0	*	*	2001	592,797	4.5	4.7	13.4	592,797	15.3	13.4			
2000	1,000,547	6.3	3.4	16.6	0	*	*	2000	631,405	4.5	4.5	13.6	631,405	15.4	13.6			
1999	1,116,976	6.1	3.8	15.7	0	*	*	1999	717,379	4.5	4.7	13.0	717,379	15.3	13.0			
1998	1,161,336	6.1	3.9	15.6	0	*	*	1998	764,296	4.7	4.8	12.6	764,296	15.1	12.6			
1997	1,144,683	6.3	4.0	15.2	0	*	*	1997	788,884	5.0	4.9	11.9	788,884	14.7	11.9			

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 20D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Summary of Trends in Growth Indicators by Age (3)														
Age and Year *	Height and Weight					Age and Year *	Height and Weight							
	Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)			Short Stature (4)	Under- weight (5)	Over- weight (5)	≥ 2 Yrs Overweight (5)				
	Number <5th	% <5th	% >95th	Number 85th-<95th	% >95th		Number 85th-<95th	% >95th	Number <5th	% <5th	% >95th	Number 85th-<95th	% >95th	
48 - 59 Months														
2006	880,596	3.8	3.6	16.4	880,596	17.1	16.4							
2005	860,503	3.8	3.8	16.3	860,503	16.8	16.3							
2004	838,883	3.9	3.9	16.1	838,883	16.7	16.1							
2003	768,211	3.9	4.2	15.9	768,211	16.4	15.9							
2002	630,982	4.0	4.5	15.7	630,982	16.2	15.7							
2001	534,308	4.0	4.3	14.4	534,308	15.7	14.4							
2000	580,987	3.9	4.2	14.2	580,987	15.9	14.2							
1999	661,574	4.0	4.4	13.5	661,574	15.6	13.5							
1998	718,459	4.2	4.4	13.0	718,459	15.6	13.0							
1997	755,563	4.4	4.5	12.4	755,563	15.3	12.4							
Total All Age Groups														
2006	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8							
2005	6,709,486	6.4	4.8	13.9	2,869,075	16.1	14.7							
2004	6,582,905	6.4	4.7	14.1	2,814,475	16.1	14.8							
2003	6,003,860	6.4	5.0	13.9	2,576,591	15.8	14.7							
2002	5,060,321	6.3	5.5	13.5	2,139,100	15.4	14.3							
2001	4,390,867	6.3	5.4	12.9	1,814,873	15.1	13.1							
2000	4,617,585	6.0	5.4	12.8	1,932,131	15.2	13.2							
1999	5,149,796	5.9	5.7	12.3	2,187,151	15.0	12.7							
1998	5,365,056	6.0	5.9	12.0	2,320,924	14.9	12.2							
1997	5,389,269	6.1	5.9	11.7	2,396,884	14.6	11.6							

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 21D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Summary of Trends in Anemia by Age (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	%
6 - 11 Months				18 - 23 Months			
	2006	456,731	16.9		2006	597,860	14.9
	2005	423,505	16.0		2005	574,477	14.4
	2004	378,164	16.5		2004	560,832	14.6
	2003	370,968	16.4		2003	463,385	14.5
	2002	383,296	16.2		2002	430,303	14.0
	2001	409,327	16.7		2001	385,770	14.3
	2000	501,041	17.2		2000	377,097	14.5
	1999	594,035	18.1		1999	430,654	15.4
	1998	596,332	18.4		1998	444,913	15.3
	1997	613,368	18.6		1997	441,511	15.3
12 - 17 Months				24 - 35 Months			
	2006	926,054	17.0		2006	976,762	15.3
	2005	869,799	16.4		2005	951,742	15.1
	2004	874,061	16.5		2004	957,443	15.0
	2003	696,251	15.8		2003	787,294	15.0
	2002	654,996	15.1		2002	748,232	14.5
	2001	581,814	15.4		2001	664,271	14.8
	2000	566,530	15.4		2000	673,890	15.1
	1999	613,229	16.7		1999	753,509	15.8
	1998	636,182	16.7		1998	800,383	15.9
	1997	611,876	16.4		1997	812,683	15.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 21D
2006 Pediatric Nutrition Surveillance (1)
National
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Summary of Trends in Anemia by Age (3)					
Age and Year		Anemia Low Hb/Hct (4)		Age and Year	
		Number	%		
				Number	%
36 - 59 Months				Total All Age Groups	
2006		1,564,184	10.3	2006	4,521,591 14.0
2005		1,561,564	10.1	2005	4,381,087 13.6
2004		1,594,401	9.9	2004	4,364,901 13.5
2003		1,282,455	9.9	2003	3,600,353 13.4
2002		1,204,360	9.6	2002	3,421,187 13.0
2001		1,075,424	9.8	2001	3,116,606 13.4
2000		1,099,010	10.0	2000	3,217,568 13.7
1999		1,254,319	10.7	1999	3,645,746 14.5
1998		1,386,489	10.9	1998	3,864,299 14.5
1997		1,445,861	11.0	1997	3,925,299 14.5
All Other/Unknown					
2006		0	*		
2005		0	*		
2004		0	*		
2003		0	*		
2002		0	*		
2001		0	*		
2000		0	*		
1999		0	*		
1998		0	*		
1997		0	*		

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

