

**STATE OF CONNECTICUT DEPARTMENT OF PUBLIC HEALTH
PRELIMINARY INFLUENZA ACTIVITY REPORT
For Week 6 (week ending February 14, 2009)**



Preliminary influenza data reviewed from the week ending February 14th revealed that a high level of activity was observed during the 20th week of the official 2008-2009 influenza season. These data include a total of 2,133 laboratory confirmed test results from various Connecticut hospitals that reveal the presence of both type A and type B flu viruses, and include reports from all 8 (Fairfield, Hartford, Litchfield, Middlesex, New Haven, New London, Tolland, and Windham) Connecticut counties (see Table 1, Figure 1). Our national flu activity classification remained as “widespread” as a result of the continued high flu activity statewide. The DPH laboratory has culture-confirmed the presence of influenza virus in 173 clinical specimens. Characterization of these isolates reveals the presence of the influenza A (H1N1) subtype in 85 specimens, the influenza A (H3N2) subtype in 35 specimens, an unspecified influenza A isolate in 1 specimen, and influenza B in 52 specimens (see Figure 2). Clinicians should be aware of oseltamivir resistance among many flu A (H1N1) strains. For the second week, influenza activity has increased to a level that exceeds weekly activity observed during the equivalent weeks in all of the last 5 seasons (see Figures 3a & 3b).

Six additional indicators of flu activity are being monitored throughout these initial weeks of the 2008-2009 flu season. Information on respiratory outbreaks and laboratory confirmed influenza cases in long-term care facilities are shown in Figure 4. Data on Connecticut residents presenting with influenza-like-illness (ILI) as reported by participants of the United States Outpatient Influenza-like Illness Surveillance Network (ILINet) (formerly known as the U.S. Influenza Sentinel Provider Surveillance Network) are shown in Figure 5. These reporting sites also include participants in the Connecticut Influenza Super Sentinel Surveillance Pilot Project organized this year to provide additional information on outpatient ILI activity.

Data on emergency department visits from our Hospital Emergency Department Syndromic Surveillance (HEDSS) System are also analyzed. One category in particular, fever/flu, has been shown to correlate well with influenza activity (see Figure 6). Patients with more severe illness who are admitted into Connecticut hospitals are tracked by the Connecticut Hospital Admissions Syndromic Surveillance (HASS) System. Data on one category, total statewide pneumonia admissions, correlates with flu activity (see Figure 7). Finally, selected mortality data are monitored each week including pneumonia and influenza deaths from Connecticut cities that participate in the U.S. 122 Cities Mortality Reporting System (see Figure 8), and reports of influenza associated pediatric deaths (none reported to date this flu season). These surveillance systems are providing additional evidence that Connecticut is currently experiencing a very high level of flu activity as compared with recent influenza seasons.

The CDC has posted its 21st national influenza activity map showing no states with initial “sporadic” flu activity, 6 states with “local” flu activity, 17 states with a higher “regional” level of activity, and 27 states with the highest “widespread” level of activity. The most recent CDC weekly influenza activity map can be viewed at: <http://www.cdc.gov/flu/weekly/usmap.htm>


If you have any questions concerning this report, please contact Alan Siniscalchi or Maria Andrews at the Department of Public Health, Epidemiology and Emerging Infections Program, by calling 860-509-7994.

**Connecticut Department of Public Health
2008-2009 Laboratory Confirmed Influenza Testing
For Week 6 (week ending February 14, 2009)**


Results by		Previous	New	Total
TEST:	Culture	36	137	173
	Rapid Test	1,306	654	1,960
FLU TYPE:	Type A	838	382	1,220
	Type B	201	124	325
	Unknown	303	285	588
COUNTY:	Fairfield	512	341	853
	Hartford	113	86	199
	Litchfield	57	28	85
	Middlesex	107	37	144
	New Haven	469	226	695
	New London	30	18	48
	Tolland	22	28	50
	Windham	32	27	59
	GENDER:	Female	682	422
	Male	660	369	1,029
Total		1,342	791	2,133

Age:	Previous	New	Total	Week	Dates - 2009	Total
0-4	111	71	182	1	January 4 - 10	40
5-24	766	523	1,289	2	January 11-17	114
25-64	395	172	567	3	January 18-24	202
≥65	70	25	95	4	January 25-31	377
Age Range: <1 – 97				5	February 1-7	591
Average Age: 22.5				6	February 8-14	720
				7	February 15-21	
				8	February 22-28	
				9	March 1-7	
				10	March 8-14	
				11	March 15-21	
				12	March 22-28	
				13	March 29- April 4	
				14	April 5-11	
				15	April 12-18	
				16	April 19-25	
				17	April 26- May 2	
				18	May 3- 9	
				19	May 10-16	
				20	May 17-23	
				21	May 24-30	
				22	May 31- June 6	
Week	Dates - 2008	Total				
40	Sept. 28 - Oct. 4	1				
41	October 5-11	1				
42	October 12-18	3				
43	October 19-25	2				
44	Oct. 26 - Nov. 1	2				
45	November 2-8	2				
46	November 9-15	2				
47	November 16-22	3				
48	November 23 -29	6				
49	Nov. 30 – Dec. 6	2				
50	December 7-13	7				
51	December 14-20	17				
52	December 21-27	19				
53	Dec. 28- Jan. 3	22				


Figure 1. Laboratory Confirmed Tests by Flu Type, Connecticut, 2008-2009 Influenza Season



**Figure 2. Department of Public Health Laboratory Culture
Confirmed Tests by Flu Type,
Connecticut, 2008-2009 Flu Season**



**Figure 3a. Laboratory Confirmed Tests by Flu Season
Connecticut, 2004-2009**



**Figure 3b. Laboratory Confirmed Tests by Flu Season
Connecticut, 2003-2009**





Figure 4. Respiratory Outbreaks & Influenza in Long-term Care Facilities (LTCF), Connecticut, 2008-2009 Influenza Season



**Figure 5. Outpatient Influenza-Like Illness Surveillance Network (ILINet),
Statewide Visits of Patients with Influenza-Like Illness (ILI), Connecticut,
2007-2009**




Figure 6. Hospital Emergency Department Syndromic Surveillance (HEDSS) System, Statewide "Fever/Flu" Visits by Flu Season, Connecticut, 2006-2009




Figure 7. Hospital Admissions Syndromic Surveillance (HASS) System, Connecticut Statewide Pneumonia Admissions by Flu Season; 2005-2009




Figure 8. CDC 122 Cities Surveillance: Pneumonia & influenza Deaths in Selected Connecticut Cities, 2007-2008 & 2008-2009 Flu Seasons

